


### Anti-ATP citrate lyase antibody [EP704Y] ab40793

KO 評価済 リコンビナント RabMAb

★★★★☆ 2 Abreviews 52 References 画像数 11

#### 製品の概要

製品名	Anti-ATP citrate lyase antibody [EP704Y]
製品の詳細	Rabbit monoclonal [EP704Y] to ATP citrate lyase
由来種	Rabbit
特異性	This antibody recognises ATP citrate lyase (ACL). The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
アプリケーション	適用あり: IHC-P, Flow Cyt (Intra), ICC/IF, WB, IP
種交差性	交差種: Mouse, Rat, Human 交差が予測される動物種: Common marmoset 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as <a href="#">ab207504</a> )
ポジティブ・コントロール	WB: HeLa cell lysate; NIH/3T3; rat lung; C6 lysates. ICC/IF: HeLa cells. Flow Cyt (intra): HeLa cells. IP: Jurkat cell lysate; HeLa. IHC: Human clear cell carcinoma of kidney
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <a href="#">see here</a> . Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> .

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS

精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP704Y
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee      Abpromise保証は、次のテスト済みアプリケーションにおけるab40793の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration.
Flow Cyt (Intra)		1/30. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. For unpurified use at 1/100.
ICC/IF		1/50.
WB	★★★★★ (2)	1/10000. Detects a band of approximately 125 kDa (predicted molecular weight: 122 kDa).Can be blocked with <b>ATP citrate lyase peptide (ab207504)</b> . <b>For unpurified use at 1/1000 - 1/5000.</b>
IP		1/20.

ターゲット情報

機能	ATP-citrate synthase is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA in many tissues. Has a central role in de novo lipid synthesis. In nervous tissue it may be involved in the biosynthesis of acetylcholine.
配列類似性	In the N-terminal section; belongs to the succinate/malate CoA ligase beta subunit family. In the C-terminal section; belongs to the succinate/malate CoA ligase alpha subunit family. Contains 1 ATP-grasp domain.
翻訳後修飾	ISGylated. Acetylated at Lys-540, Lys-546 and Lys-554 by KAT2B/PCAF. Acetylation is promoted by glucose and stabilizes the protein, probably by preventing ubiquitination at the same sites. Acetylation promotes de novo lipid synthesis. Deacetylated by SIRT2. Ubiquitinated at Lys-540, Lys-546 and Lys-554 by UBR4, leading to its degradation. Ubiquitination is probably inhibited by acetylation at same site.
細胞内局在	Cytoplasm.

画像



Western blot - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)


**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)

**Lane 2:** ATP citrate lyase knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** HeLa whole cell lysate (20 µg)

**Lanes 1 - 3:** Merged signal (red and green). Green - ab40793 observed at 125 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

Unpurified ab40793 was shown to specifically react with ATP citrate lyase in wild-type HAP1 cells as signal was lost in ATP citrate lyase knockout cells. Wild-type and ATP citrate lyase knockout samples were subjected to SDS-PAGE. ab40793 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

Immunocytochemistry/Immunofluorescence analysis of HeLa cells labelling ATP citrate lyase with purified ab40793 at 1/50. Cells were fixed with 100% methanol. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody.


Control: PBS only.

Nuclear counter stain: DAPI.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human clear cell carcinoma of kidney tissue sections labeling ATP citrate lyase with Purified ab40793 at 1:100 dilution. Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry (Intracellular) - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling ATP citrate lyase with purified ab40793 at 1/30 dilution (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

**All lanes :** Anti-ATP citrate lyase antibody [EP704Y] (ab40793) at 1/50000 dilution (purified)

**Lane 1 :** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

**Lane 2 :** NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates


Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

**Predicted band size:** 122 kDa

Blocking and diluting buffer: 5% NFDM/TBST.



Immunoprecipitation - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

ab40793 (purified) at 1:20 dilution (1.5µg) immunoprecipitating ATP citrate lyase in HeLa whole cell lysate.


**Lane 1 (input):** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate, 10µg

**Lane 2 (+):** ab40793 & HeLa whole cell lysate

**Lane 3 (-):** Rabbit monoclonal IgG (**ab172730**) instead of ab40793 in HeLa whole cell lysate.

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST.



**All lanes :** Anti-ATP citrate lyase antibody [EP704Y] (ab40793) at 1/10000 dilution (purified)

**Lane 1 :** Rat lung lysates

**Lane 2 :** C6 (Rat glial tumor glial cell) whole cell lysate


Lysates/proteins at 15 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 122 kDa


Blocking and diluting buffer: 5% NFDM/TBST.



Anti-ATP citrate lyase antibody [EP704Y] (ab40793) at 1/5000 dilution (unpurified) + HeLa cell lysate


**Predicted band size:** 122 kDa

**Observed band size:** 122 kDa



Flow Cytometry (Intracellular) - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

Overlay histogram showing HeLa cells stained with unpurified ab40793 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab40793, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1 µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed.



Immunoprecipitation - Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

Unpurified ab40793 at 1/40 immunoprecipitating ATP citrate lyase in HeLa (human cervix adenocarcinoma) whole cell lysate.

Lane 1 (input): HeLa (human cervix adenocarcinoma) whole cell lysate 10 µg

Lane 2 (+): ab40793 + HeLa (human cervix adenocarcinoma) whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab40793 in HeLa (human cervix adenocarcinoma) whole cell lysate

For western blotting, ab40793 at 1/1000 dilution and [ab131366](#) VeriBlot for IP (HRP) was used for detection at 1/10000.

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-ATP citrate lyase antibody [EP704Y] (ab40793)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors