


# Anti-ATPB antibody [3D5] - Mitochondrial Marker ab14730

★★★★★ **41 Abreviews**   **229 References**   画像数 6

### 製品の概要

製品名	Anti-ATPB antibody [3D5] - Mitochondrial Marker
製品の詳細	Mouse monoclonal [3D5] to ATPB - Mitochondrial Marker
由来種	Mouse
特異性	Human and Bovine complex V beta subunit (ATPB).
アプリケーション	<b>適用あり:</b> WB, ICC/IF, Flow Cyt, IHC-P
種交差性	<b>交差種:</b> Mouse, Rat, Cow, Human, Caenorhabditis elegans <b>交差が予測される動物種:</b> Goat, Cat, Dog, Pig, Common marmoset 
免疫原	Tissue, cells or virus corresponding to Human ATPB.
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.4 Preservative: 0.02% Sodium azide Constituent: HEPES buffered saline
精製度	IgG fraction
特記事項 (精製)	Near homogeneity as judged by SDS-PAGE. The antibody was produced in vitro using hybridomas grown in serum-free medium, and then purified by biochemical fractionation.
ポリ/モノ	モノクローナル

クローン名	3D5
アイソタイプ	IgG1
軽鎖の種類	kappa

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab14730の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (14)	Use a concentration of 0.5 µg/ml. Detects a band of approximately 52 kDa (predicted molecular weight: 52 kDa).
ICC/IF	★★★★★ (8)	Use a concentration of 1 - 2 µg/ml.
Flow Cyt		Use a concentration of 1 µg/ml. <b>ab170190</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IHC-P	★★★★★ (10)	Use at an assay dependent concentration.


## ターゲット情報

**機能**      Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Subunits alpha and beta form the catalytic core in F(1). Rotation of the central stalk against the surrounding alpha(3)beta(3) subunits leads to hydrolysis of ATP in three separate catalytic sites on the beta subunits.

**配列類似性**      Belongs to the ATPase alpha/beta chains family.

**細胞内局在**      Mitochondrion. Mitochondrion inner membrane. Peripheral membrane protein.

## 画像



Western blot - Anti-ATPB antibody [3D5] -  
Mitochondrial Marker (ab14730)

**All lanes :**

**Lane 1 :** HeLa

**Lane 2 :** HepG2

**Lane 3 :** Hu liver

**Lane 4 :** Ms liver


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 52 kDa


Merged signal (red and green). Green - ab14730 observed at 52 kDa. Red - loading control, **ab181602** observed at 37 kDa.

Samples were subjected to SDS-PAGE. ab14730 and **ab181602** (Rabbit anti-GAPDH loading control) were incubated overnight at 4°C at 1 µg/mL and 1/10000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed (**ab216772**) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed (**ab216777**) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730)

ab14730 staining ATPB in cultured human fibroblasts. Cells were fixed, permeabilized and then labelled with ab14730 followed by an AlexaFluor® 488-conjugated Goat anti-Mouse IgG1-specific secondary antibody (2 µg/ml)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730)

ab14730 (2µg/ml) staining ATPB in human duodenum using an automated system (DAKO Autostainer Plus). Using this protocol there is cytoplasmic and mitochondrial staining of epithelium. Sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffer EDTA pH 9.0 in a DAKO PT link. Slides were peroxidase blocked in 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that, for manual staining, optimization of primary antibody concentration and incubation time is recommended. Signal amplification may be required.



Flow Cytometry - Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730)

ab14730 (blue) at 1µg/ml staining ATPB in HL-60 cells and analyzed by Flow cytometry. Red histogram represents equal quantity of isotype control.



Western blot - Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730)

**All lanes :** Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730)

**Lane 1 :** isolated mitochondria from human heart at 5 µg


**Lane 2 :** isolated mitochondria from bovine heart at 1 µg

**Lane 3 :** isolated mitochondria from rat heart at 10 µg

**Lane 4 :** isolated mitochondria from mouse heart at 10 µg

**Predicted band size:** 52 kDa

**Observed band size:** 52 kDa



Western blot - Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730)

**All lanes :** Anti-ATPB antibody [3D5] - Mitochondrial Marker (ab14730) at 0.8 µg/ml

**Lane 1 :** HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

**Lane 2 :** HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

**Lane 3 :** Human liver tissue lysate - total protein ([ab29889](#))

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

**Predicted band size:** 52 kDa

**Observed band size:** 52 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors