


Anti-ATXN2L antibody ab99304

画像数 2

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-ATXN2L antibody |
| 製品の詳細 | Rabbit polyclonal to ATXN2L |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: WB, IP |
| 種交差性 | 交差種: Human 交差が予測される動物種: Mouse, Rat, Rabbit, Guinea pig, Dog, Pig, Chimpanzee, Rhesus monkey, Gorilla, Orangutan, Elephant  |
| 免疫原 | Synthetic peptide within Human ATXN2L aa 150-200. The exact sequence is proprietary. Database link: NP_009176.2 |
| ポジティブ・コントロール | WB: HeLa and 293T whole cell lysate. |
| 特記事項 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|--------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| バッファー | pH: 6.8 Preservative: 0.09% Sodium azide Constituents: Tris buffered saline, 0.1% BSA |
| 精製度 | Immunogen affinity purified |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab99304の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| WB | | 1/2000 - 1/10000. Predicted molecular weight: 114 kDa. |
| IP | | Use at 2-5 µg/mg of lysate. |

ターゲット情報

| | |
|-------|--|
| 組織特異性 | Expressed at high levels in thymus, lymph node, spleen, fetal kidney and adult testis. Constitutively associated with MPL and EPOR in hematopoietic cells. |
| 配列類似性 | Belongs to the ataxin-2 family. |
| 翻訳後修飾 | Thrombopoietin triggers the phosphorylation on tyrosine residues in a way that is dependent on MPL C-terminal domain. Arg-361 is dimethylated, probably to asymmetric dimethylarginine. |
| 細胞内局在 | Membrane. |

画像




All lanes : Anti-ATXN2L antibody (ab99304) at 0.04 µg/ml

- Lane 1 :** Hela whole cell lysate at 50 µg
- Lane 2 :** Hela whole cell lysate at 15 µg
- Lane 3 :** Hela whole cell lysate at 5 µg
- Lane 4 :** 293T whole cell lysate at 50 µg

Developed using the ECL technique.

Predicted band size: 114 kDa

Exposure time: 30 seconds



ab99304, at 0.1 µg/ml, staining ATXN2L in HeLa whole cell lysate immunoprecipitated using ab99304 at 3 µg/mg lysate (1 mg/IP; 20% of IP loaded/lane). Detection: Chemiluminescence with an exposure time of 30 seconds.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors