


### Anti-BAIAP2L1/IRTKS antibody ab96349

画像検索

医薬用外劇物

#### 製品の概要

製品名	Anti-BAIAP2L1/IRTKS antibody
製品の詳細	Rabbit polyclonal to BAIAP2L1/IRTKS
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, WB
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat 
免疫原	Recombinant fragment corresponding to Human BAIAP2L1/IRTKS aa 1-263.
ポジティブ・コントロール	293T, A431, HeLa or HepG2 cell lysate. ICC/IF: A431 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.00 Preservative: 0.01% Thimerosal (merthiolate) Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

## アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab96349の使用に適用されず**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/100 - 1/1000.
WB		1/500 - 1/3000. Predicted molecular weight: 57 kDa.

## ターゲット情報


機能	May function as adapter protein. Involved in the formation of clusters of actin bundles. Plays a role in the reorganization of the actin cytoskeleton in response to bacterial infection.
配列類似性	Contains 1 IMD (IRSp53/MIM homology) domain. Contains 1 SH3 domain.
ドメイン	The IMD domain is predicted to have a helical structure. It may induce actin bundling and filopodia formation.
翻訳後修飾	Phosphorylated on tyrosine in response to insulin.
細胞内局在	Cytoplasm, cytoskeleton. Recruited to actin pedestals that are formed upon infection by bacteria at bacterial attachment sites.

## 画像



Anti-BAIAP2L1/IRTKS antibody (ab96349) at 1/1000 dilution +  
HeLa whole cell lysate at 30 µg

**Predicted band size: 57 kDa**



Methanol-fixed A431 cell labeling BAIAP2L1/IRTKS using ab96349 at 1/200 dilution (green) in ICC/IF.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors