

# Anti-BCRP/ABCG2 antibody [BXP-21] ab3380

★★★★☆ [11 Abreviews](#) [113 References](#)

### 製品の概要

製品名	Anti-BCRP/ABCG2 antibody [BXP-21]
製品の詳細	Mouse monoclonal [BXP-21] to BCRP/ABCG2
由来種	Mouse
特異性	BXP-21 reacts with an internal epitope of BCRP/ABCG2, a 70 kD transmembrane half-transporter which is involved in Multidrug resistance. This antibody does not react with the human MDR1, MRP1 and MRP2 gene products.
アプリケーション	<b>適用あり:</b> IHC-Fr, IHC-P, ICC/IF, Flow Cyt, ICC
種交差性	<b>交差種:</b> Human
免疫原	Recombinant fragment within BCRP/ABCG2 aa 250-400. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please <b>contact</b> our Scientific Support team to discuss your requirements.
ポジティブ・コントロール	IHC-P: Human breast cancer tissue. ICC/IF: MCF7 cells. Caco Cell lysate (see second reference cited).
特記事項	<p>For the WB, please note the subcellular localization and distribution of the BCRP/ABCG2 in Normal Human Tissues: PMID:11309308</p> <p>We have received mixed feedback from customers, the antibody showed 45-50 kDa band size in A549 and Caco2 cell lines against predicted size of 72kDa. In response to data we have received we no longer guarantee this antibody for western blot. ABCG2 is a multi-pass membrane protein so for troubleshooting we would recommend heating the samples at 60-70C for 15-20 minutes instead of boiling and as this protein exists as homodimer so complete reduction of samples is highly recommended. In case of any other question please email scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

## 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.3 Constituent: 0.7% BSA
精製度	Tissue culture supernatant
一次抗体 備考	Breast Cancer Resistance Protein (BCRP) is a 70 kDa ATP-Binding Cassette membrane transport protein involved in multidrug resistance. BCRP may be over-expressed in cancer cell lines selected with doxorubicin / verapamil, topotecan or mitoxantrone.
ポリ/モノ	モノクローナル
クローン名	BXP-21
アイソタイプ	IgG2a

## アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab3380の使用に適用されます**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr	★★★★★ (1)	Use at an assay dependent concentration.
IHC-P	★★★★★ (2)	Use at an assay dependent concentration.
ICC/IF	★★★★★ (2)	Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. PubMed: 24136221 <b>ab170191</b> - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
ICC		Use at an assay dependent concentration. Acetone fixed cells.

## ターゲット情報

機能	Xenobiotic transporter that may play an important role in the exclusion of xenobiotics from the brain. May be involved in brain-to-blood efflux. Appears to play a major role in the multidrug resistance phenotype of several cancer cell lines. When overexpressed, the transfected cells become resistant to mitoxantrone, daunorubicin and doxorubicin, display diminished intracellular accumulation of daunorubicin, and manifest an ATP-dependent increase in the efflux of rhodamine 123.
組織特異性	Highly expressed in placenta. Low expression in small intestine, liver and colon.
配列類似性	Belongs to the ABC transporter superfamily. ABCG family. Eye pigment precursor importer (TC 3.A.1.204) subfamily. Contains 1 ABC transmembrane type-2 domain. Contains 1 ABC transporter domain.

翻訳後修飾

Glycosylation-deficient ABCG2 is normally expressed and functional.

細胞内局在

Cell membrane.

---

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors