


Anti-Bestrophin/BEST1 antibody [E6-6] ab2182

★★★★★ [7 Abreviews](#) [30 References](#) [画像数 2](#)

製品の概要

製品名	Anti-Bestrophin/BEST1 antibody [E6-6]
製品の詳細	Mouse monoclonal [E6-6] to Bestrophin/BEST1
由来種	Mouse
アプリケーション	適用あり: WB, IHC-Fr, ICC/IF, IP
種交差性	交差種: Cow, Dog, Human, Pig, Monkey 交差が予測される動物種: Non human primates  非交差種: Mouse, Rat, Rabbit, Goat
免疫原	Synthetic peptide corresponding to Human Bestrophin/BEST1 aa 568-585 (C terminal) conjugated to keyhole limpet haemocyanin. Sequence: KDHMDPYWALENRDEAHS Database link: O76090 Run BLAST with Run BLAST with
ポジティブ・コントロール	IHC-Fr: Pig retinal pigment epithelium tissue. ICC/IF: Bovine retinal pigment epithelium (RPE). WB: Human RPE cell lysate.
特記事項	This product has switched from ascites to TCS on 9th September 2020. The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide

	Constituent: 99% PBS
精製度	Protein A/G purified
ポリ/モノ	モノクローナル
クローン名	E6-6
ミエローマ	unknown
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab2182の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (3)	1/1000.
IHC-Fr	★★★★★ (2)	Use at an assay dependent concentration.
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration.
IP	★★★★★ (1)	Use at an assay dependent concentration.

ターゲット情報

機能	Forms calcium-sensitive chloride channels. Highly permeable to bicarbonate.
組織特異性	Predominantly expressed in the basolateral membrane of the retinal pigment epithelium.
関連疾患	<p>Defects in BEST1 are the cause of vitelliform macular dystrophy type 2 (VMD2) [MIM:153700]; also known as Best macular dystrophy (BMD). VMD2 is an autosomal dominant form of macular degeneration that usually begins in childhood or adolescence. VMD2 is characterized by typical 'egg-yolk' macular lesions due to abnormal accumulation of lipofuscin within and beneath the retinal pigment epithelium cells. Progression of the disease leads to destruction of the retinal pigment epithelium and vision loss.</p> <p>Defects in BEST1 are the cause of retinitis pigmentosa type 50 (RP50) [MIM:613194]. A retinal dystrophy belonging to the group of pigmentary retinopathies. RP is characterized by retinal pigment deposits visible on fundus examination and primary loss of rod photoreceptor cells followed by secondary loss of cone photoreceptors. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well.</p> <p>Defects in BEST1 are a cause of adult-onset vitelliform macular dystrophy (AVMD) [MIM:608161]. AVMD is a rare autosomal dominant disorder with incomplete penetrance and highly variable expression. Patients usually become symptomatic in the fourth or fifth decade of life with a protracted disease of decreased visual acuity.</p> <p>Defects in BEST1 are the cause of bestrophinopathy autosomal recessive (ARB) [MIM:611809]. A retinopathy characterized by central visual loss, an absent electro-oculogram light rise, and a reduced electroretinogram.</p> <p>Defects in BEST1 are the cause of vitreoretinchoroidopathy autosomal dominant (ADVIRC)</p>

[MIM:193220]. A disorder characterized by vitreoretinchoroidal dystrophy. The clinical presentation is variable and may be associated with cataract, nanophthalmos, microcornea, shallow anterior chamber, and glaucoma.

配列類似性

Belongs to the bestrophin family.


翻訳後修飾

Phosphorylated by PP2A.

細胞内局在


Cell membrane. Basolateral cell membrane.

画像



Western blot - Anti-Bestrophin/BEST1 antibody [E6-6] (ab2182)

Anti-Bestrophin/BEST1 antibody [E6-6] (ab2182) at 1/1000 dilution
+ Human RPE cell lysate



Western blot - Anti-Bestrophin/BEST1 antibody [E6-6] (ab2182)

This image is courtesy of an Abreview submitted by Dr Madimir Milenkovic

All lanes : Anti-Bestrophin/BEST1 antibody [E6-6] (ab2182) at 1/1000 dilution

Lane 1 : Human RPE, retinal pigment epithelial cell lysate

Lane 2 : Non transfected HEK 293 cell extract

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : HRP-conjugated goat anti-mouse

Developed using the ECL technique.

Performed under reducing conditions.

Observed band size: 67 kDa

Exposure time: 5 minutes

The primary antibody was diluted in PBS/Tween/5%Milk and incubated for 1.5 hours at 25°C.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors