

# Biotin Anti-CCR7 antibody [4B12] ab93519

### 3 References

#### 製品の概要

---

|              |   |
|--------------|---|
| 製品名          | Biotin Anti-CCR7 antibody [4B12]  |
| 製品の詳細        | Biotin Rat monoclonal [4B12] to CCR7  |
| 由来種          | Rat   |
| 標識           | Biotin  |
| アプリケーション     | <b>適用あり:</b> Flow Cyt, ICC/IF   |
| 種交差性         | <b>交差種:</b> Mouse   |
| 免疫原          | Tissue, cells or virus corresponding to Mouse CCR7. Rat basophilic leukaemia (RBL)-2H3 cells transfected with Mouse CCR7.   |
| ポジティブ・コントロール | Splenocytes or thymocytes.  |
| 特記事項         | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

#### 製品の特性

---

|        |   |
|--------|---|
| 製品の状態  | Liquid  |
| 保存方法   | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー  | pH: 7.20<br>Preservative: 0.09% Sodium azide<br>Constituents: 0.87% Sodium chloride, PBS                  |
| 精製度    | Protein G purified  |
| ポリ/モノ  | モノクローナル   |
| クローン名  | 4B12  |
| アイソタイプ | IgG2a   |
| 軽鎖の種類  | kappa   |

## アプリケーション

---

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab93519の使用に適用されます**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項                                     |
|----------|-----------|--|
| Flow Cyt |           | Use at an assay dependent concentration. |
| ICC/IF   |           | Use at an assay dependent concentration. |

## ターゲット情報

---

|       |   |
|-------|---|
| 機能    | Receptor for the MIP-3-beta chemokine. Probable mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions.   |
| 組織特異性 | Expressed in various lymphoid tissues and activated B- and T-lymphocytes, strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7. |
| 配列類似性 | Belongs to the G-protein coupled receptor 1 family.   |
| 細胞内局在 | Cell membrane.  |

---

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors