

Anti-BubR1 (phospho T680) antibody [EPR19958] ab200061

リコンビナント RabMAb[®]

1 References [画像数 5](#)

製品の概要

| | |
|--------------|--|
| 製品名 | Anti-BubR1 (phospho T680) antibody [EPR19958] |
| 製品の詳細 | Rabbit monoclonal [EPR19958] to BubR1 (phospho T680) |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: IP, ICC/IF, WB, Flow Cyt (Intra) |
| 種交差性 | 交差種: Human |
| 免疫原 | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. |
| ポジティブ・コントロール | WB: HeLa whole cell lysate treated with 0.5 μ M nocodazole for 24 hours. ICC/IF: HeLa cells. Flow Cyt (intra): HeLa (human cervix adenocarcinoma) treated with 0.5 ng/ml nocodazole for 24 hours. IP: HeLa whole cell lysate treated with 0.5 μ M nocodazole for 24 hours. |
| 特記事項 | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> |

製品の特性

| | |
|-------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |
| バッファー | pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| 精製度 | Protein A purified |
| ポリ/モノ | モノクローナル |
| クローン名 | EPR19958 |

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab200061の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|------------------|-----------|--|
| IP | | 1/30. |
| ICC/IF | | 1/100. |
| WB | | 1/2000. Detects a band of approximately 120 kDa (predicted molecular weight: 120 kDa). |
| Flow Cyt (Intra) | | 1/500. |

ターゲット情報

機能

Essential component of the mitotic checkpoint. Required for normal mitosis progression. The mitotic checkpoint delays anaphase until all chromosomes are properly attached to the mitotic spindle. One of its checkpoint functions is to inhibit the activity of the anaphase-promoting complex/cyclosome (APC/C) by blocking the binding of CDC20 to APC/C, independently of its kinase activity. The other is to monitor kinetochore activities that depend on the kinetochore motor CENPE. Required for kinetochore localization of CENPE. Negatively regulates PLK1 activity in interphase cells and suppresses centrosome amplification. Also implicated in triggering apoptosis in polyploid cells that exit aberrantly from mitotic arrest. May play a role for tumor suppression.

組織特異性

Highly expressed in thymus followed by spleen. Preferentially expressed in tissues with a high mitotic index.

関連疾患

Note=Defects in BUB1B are associated with tumor formation.

Defects in BUB1B are the cause of premature chromatid separation trait (PCS) [MIM:176430]. PCS consists of separate and splayed chromatids with discernible centromeres and involves all or most chromosomes of a metaphase. It is found in up to 2% of metaphases in cultured lymphocytes from approximately 40% of normal individuals. When PCS is present in 5% or more of cells, it is known as the heterozygous PCS trait and has no obvious phenotypic effect, although some have reported decreased fertility. Inheritance is autosomal dominant.

Defects in BUB1B are the cause of mosaic variegated aneuploidy syndrome (MVA) [MIM:257300]. MVA is a severe autosomal recessive developmental disorder characterized by mosaic aneuploidies, predominantly trisomies and monosomies, involving multiple different chromosomes and tissues. The proportion of aneuploid cells varies but is usually more than 25% and is substantially greater than in normal individuals. Affected individuals typically present with severe intrauterine growth retardation and microcephaly. Eye anomalies, mild dysmorphism, variable developmental delay, and a broad spectrum of additional congenital abnormalities and medical conditions may also occur. The risk of malignancy is high, with rhabdomyosarcoma, Wilms tumor and leukemia reported in several cases. MVA is caused by biallelic mutations in the BUB1B gene.

配列類似性

Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. BUB1 subfamily.
Contains 1 BUB1 N-terminal domain.
Contains 1 protein kinase domain.

ドメイン

The D-box targets the protein for rapid degradation by ubiquitin-dependent proteolysis during the transition from mitosis to interphase.

The BUB1 N-terminal domain directs kinetochore localization and binding to BUB3.

翻訳後修飾

Proteolytically cleaved by caspase-3 in a cell cycle specific manner. The cleavage might be involved in the durability of the cell cycle delay. Caspase-3 cleavage is associated with abrogation of the mitotic checkpoint. The major site of cleavage is at Asp-610.

Acetylation at Lys-250 regulates its degradation and timing in anaphase entry.

Ubiquitinated. Degradated by the proteasome.

Sumoylated by SUMO2 and SUMO3. The sumoylation mediates the association with CENPE at the kinetochore.

Autophosphorylated in vitro. Intramolecular autophosphorylation is stimulated by CENPE.


Phosphorylated during mitosis and hyperphosphorylated in mitotically arrested cells.

Phosphorylation at Ser-670 and Ser-1043 occurs at kinetochores upon mitotic entry with dephosphorylation at the onset of anaphase.

細胞内局在

Cytoplasm. Nucleus. Chromosome > centromere > kinetochore. Cytoplasm > cytoskeleton > centrosome. Cytoplasmic in interphase cells. Associates with the kinetochores in early prophase. Kinetochore localization requires BUB1, PLK1 and CASC5.

画像



Immunocytochemistry/ Immunofluorescence - Anti-BubR1 (phospho T680) antibody [EPR19958] (ab200061)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling BubR1 (phospho T680) with ab200061 at 1/100 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green).

Confocal image showing positive staining on M phase HeLa cells, and LP treatment completely blocked the staining. (PMID:11792804).

The nuclear counter stain is DAPI (blue).

Tubulin is detected with [ab195889](#) (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)) at 1/250 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is [ab150077](#) at 1/1000 dilution.



All lanes : Anti-BubR1 (phospho T680) antibody [EPR19958] (ab200061) at 1/2000 dilution

Lane 1 : Untreated HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 2 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate treated with 0.5 μ M nocodazole for 24 hours

Lane 3 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate treated with 0.5 μ M nocodazole for 24 hours, then treated with FastAP Thermosensitive Alkaline Phosphatase for 1 hour

Lysates/proteins at 10 μ g per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Predicted band size: 120 kDa

Observed band size: 120 kDa


Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.



Flow Cytometry (Intracellular) - Anti-BubR1 (phospho T680) antibody [EPR19958] (ab200061)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed HeLa (Human epithelial cell line from cervix adenocarcinoma) cells, untreated or treated with 0.5 ng/ml nocodazole for 24 hours, labeling BubR1 (phospho T680) with ab200061 at 1/500 dilution compared with a Rabbit IgG, monoclonal [EPR25A]-Isotype control (**ab172730**) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat anti Rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-BubR1 (phospho T680) antibody [EPR19958] (ab200061)

BubR1 (phospho T680) was immunoprecipitated from 0.35 mg of HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate treated with 0.5 μ M nocodazole for 24 hours with ab200061 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab200061 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: HeLa whole cell lysate treated with 0.5 μ M nocodazole for 24 hours 10 μ g (Input).

Lane 2: ab200061 IP in HeLa whole cell lysate treated with 0.5 μ M nocodazole for 24 hours.

Lane 3: Rabbit IgG, monoclonal [EPR25A]-Isotype Control (**ab172730**) instead of ab200061 in HeLa whole cell lysate treated with 0.5 μ M nocodazole for 24 hours.

Blocking and dilution buffer and concentration: 5% NFDN/TBST.

Exposure time: 3 seconds.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-BubR1 (phospho T680) antibody [EPR19958]
(ab200061)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors