



### Anti-Caspase-3 antibody ab49822

★★★★★ [7 Abreviews](#) [388 References](#) [画像数 2](#)

#### 製品の概要

製品名	Anti-Caspase-3 antibody
製品の詳細	Rabbit polyclonal to Caspase-3
由来種	Rabbit
特異性	<p>The immunogen for this product is aa167-175 within the p17 subunit (cleaved caspase) but because this is a poly the epitope will vary and not all batches will be flush against the cleavage site.</p> <p>This means it will detect a number of additional cleavage products in addition to p17:</p> <ul style="list-style-type: none"><li>• P12+17,</li><li>• P12+P17+propeptide 1</li><li>• p12+p17+propeptide 1 + propeptide 2,</li><li>• P17+propeptide 1,</li><li>• p17+propeptide 1 + propeptide 2</li></ul>
アプリケーション	<b>適用あり:</b> WB, ICC/IF
種交差性	<b>交差種:</b> Human
免疫原	<p>Synthetic peptide corresponding to Human Caspase-3 aa 150-250 conjugated to keyhole limpet haemocyanin.</p> <p>Database link: <a href="#">P42574</a></p> <p> <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a></p>
ポジティブ・コントロール	ICC/IF: Human epitheloid carcinoma HeLa cell line, treated with staurosporine and WB: Human recombinant caspase-3, active.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

#### バッファー

pH: 7.40  
Preservative: 0.097% Sodium azide  
Constituent: 0.0268% PBS

#### 精製度

IgG fraction

#### 特記事項 (精製)

Whole antiserum is fractionated and then further purified by ion exchange chromatography. The resulting IgG fraction is further purified by absorption on the Caspase-3 peptide (Human, amino acids 167-178) corresponding to the uncleaved Caspase 3 site sequence.

#### ポリモノ

ポリクローナル

#### アイソタイプ

IgG

### アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab49822の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★☆ (2)	1/500.
ICC/IF		1/1000.

### ターゲット情報

#### 機能

Involved in the activation cascade of caspases responsible for apoptosis execution. At the onset of apoptosis it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7 and -9. Involved in the cleavage of huntingtin.

#### 組織特異性

Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.

#### 配列類似性

Belongs to the peptidase C14A family.


#### 翻訳後修飾

Cleavage by granzyme B, caspase-6, caspase-8 and caspase-10 generates the two active subunits. Additional processing of the propeptides is likely due to the autocatalytic activity of the activated protease. Active heterodimers between the small subunit of caspase-7 protease and the large subunit of caspase-3 also occur and vice versa.  
S-nitrosylated on its catalytic site cysteine in unstimulated human cell lines and denitrosylated upon activation of the Fas apoptotic pathway, associated with an increase in intracellular caspase activity. Fas therefore activates caspase-3 not only by inducing the cleavage of the caspase zymogen to its active subunits, but also by stimulating the denitrosylation of its active site thiol.

#### 細胞内局在

Cytoplasm.

### 画像



Western blot - Anti-Caspase-3 antibody (ab49822)


Anti-Caspase-3 antibody (ab49822) at 1/1000 dilution

**Secondary**

Goat Anti-Rabbit IgG, Peroxidase conjugate

Developed using the ECL technique.

**Observed band size:** 17 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Caspase-3 antibody (ab49822)

Immunocytochemistry/ Immunofluorescence analysis of HeLa cells labeling Cleaved Caspase-3 with ab49822 at 1/1000. HeLa cells were treated with 1 µg/mL staurosporine, fixed and permeabilized with Methanol followed by Acetone. Goat Anti-Rabbit IgG (whole molecule)-FITC was used as the secondary antibody. Cells were counterstained with DAPI (blue) to stain nuclei.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors