




# Anti-Caveolin-1 antibody - Caveolae Marker ab2910

★★★★★ [54 Abreviews](#) [174 References](#) [画像数 9](#)

### 製品の概要

製品名	Anti-Caveolin-1 antibody - Caveolae Marker
製品の詳細	Rabbit polyclonal to Caveolin-1 - Caveolae Marker
由来種	Rabbit
アプリケーション	<b>適用あり:</b> WB, ICC/IF, IP
種交差性	<b>交差種:</b> Mouse, Rat, Human <b>交差が予測される動物種:</b> Sheep, Rabbit, Horse, Cow, Cat, Pig, Chimpanzee, Gorilla, African green monkey, African bush elephant 
免疫原	Synthetic peptide corresponding to Human Caveolin-1 aa 1-100. Database link: <a href="#">Q03135</a> <div>  <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a> </div>
ポジティブ・コントロール	WB: Human lung, heart and spleen tissue lysates; Mouse heart and lung tissue lysates; Rat heart protein extract; PANC-1, U-87 MG, A549, HeLa, PC-3, U-2 OS, C2C12 and A431 cell lysates. ICC/IF: A-375, HeLa and NIH/3T3 cells; Rat astrocytes.
特記事項	This antibody can be used as a marker for lipid raft fractions.  The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.  If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル

## アプリケーション


**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab2910の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (21)	Use a concentration of 1 - 2 µg/ml. Detects a band of approximately 22 kDa (predicted molecular weight: 20 kDa).
ICC/IF	★★★★★ (15)	Use at an assay dependent concentration.
IP	★★★★★ (9)	Use at an assay dependent concentration. Recommended dilution: 5 µg

## ターゲット情報

機能	May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein alpha subunits and can functionally regulate their activity (By similarity). Involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Recruits CTNNB1 to caveolar membranes and may regulate CTNNB1-mediated signaling through the Wnt pathway.
組織特異性	Expressed in muscle and lung, less so in liver, brain and kidney.
関連疾患	Defects in CAV1 are the cause of congenital generalized lipodystrophy type 3 (CGL3) [MIM:612526]; also called Berardinelli-Seip congenital lipodystrophy type 3 (BSCL3). Congenital generalized lipodystrophies are autosomal recessive disorders characterized by a near absence of adipose tissue, extreme insulin resistance, hypertriglyceridemia, hepatic steatosis and early onset of diabetes.
配列類似性	Belongs to the caveolin family.
翻訳後修飾	The initiator methionine for isoform Beta is removed during or just after translation. The new N-terminal amino acid is then N-acetylated.
細胞内局在	Golgi apparatus membrane. Cell membrane. Membrane > caveola. Membrane raft. Colocalized with DPP4 in membrane rafts. Potential hairpin-like structure in the membrane. Membrane protein of caveolae.

## 画像



**All lanes** : Anti-Caveolin-1 antibody - Caveolae Marker (ab2910) at 1.5 µg/ml

**Lane 1** : Human lung

**Lane 2** : Human heart

**Lane 3** : Human spleen


Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes** : Alexa Fluor anti-rabbit at 1/5000 dilution


**Predicted band size:** 20 kDa

**Observed band size:** 20 kDa



ab2910 staining Caveolin-1 in wild-type HeLa cells (top panel) and CAV1 knockout HeLa cells ([ab255371](#)) (bottom panel). The cells were fixed with 100% methanol (5 min) then permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab2910 at 1/500 dilution and [ab7291](#) (Mouse monoclonal to alpha Tubulin) at 1/1000 dilution overnight at 4°C followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) ([ab150081](#)) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) ([ab150120](#)) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems TCS SP8).




Caveolin 1 was immunoprecipitated using 5 µg of ab2910 from lysate of Mouse Heart (Lane 3) using the Dynabeads® Protein A Immunoprecipitation Kit. Normal Rabbit IgG was used as a Isotype control (Lane 2). 10 % input represents the cell extract used for immunoprecipitation (Lane 1). Western blot analysis was performed using Caveolin 1 Rabbit Polyclonal Antibody and Goat anti-Rabbit IgG (H+L) Superclonal™ Secondary Antibody, HRP conjugate. Chemiluminescent detection was performed using Pierce™ ECL Western Blotting Substrate.

**All lanes :**

**Lane 1 :** Input control

**Lane 2 :** Isotype control

**Lane 3 :** IP elute



**All lanes :** Anti-Caveolin-1 antibody - Caveolae Marker (ab2910) at 1 µg/ml


**Lane 1 :** CAV1 knockout HeLa cell lysate

**Lane 2 :** Wild-type HeLa cell lysate

**Predicted band size:** 20 kDa

The specificity of ab2910 was demonstrated by CRISPR targeted CAV1 knockout in HeLa cells. Western blot analysis of whole cell lysates using this antibody showed no detection of caveolin 1 protein expression in knockout cells compared to the protein detected at ~22kDa in wild-type HeLa cells.

**Knockout validation info.**



**All lanes :** Anti-Caveolin-1 antibody - Caveolae Marker (ab2910) at 1 µg/ml

**Lane 1 :** PANC-1 (Human pancreatic epithelial carcinoma cell line) whole cell lysate

**Lane 2 :** U-87 MG (Human glioblastoma-astrocytoma epithelial cell line) whole cell lysate

**Lane 3 :** A549 (Human lung carcinoma cell line) whole cell lysate

**Lane 4 :** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

**Lane 5 :** PC-3 (Human prostate adenocarcinoma cell line) whole cell lysate

**Lane 6 :** U-2 OS (Human bone osteosarcoma epithelial cell line) whole cell lysate

**Lane 7 :** Mouse heart tissue lysate

**Lane 8 :** Rat heart tissue lysate

**Lane 9 :** C2C12 (Mouse myoblast cell line) whole cell lysate


**Lane 10 :** Mouse lung tissue lysate

**Lane 11 :** A-431 (Human epidermoid carcinoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.


**Predicted band size:** 20 kDa

Western blot analysis was performed on whole cell extracts (20 µg lysate). The blots were probed with Anti-ab2910 (1-2 µg/mL) and detected by chemiluminescence using Goat anti-Rabbit IgG (H+L) Superclonal™ Secondary Antibody, HRP conjugate. A 17 kDa band corresponding to Caveolin-1 was observed across cell lines and tissues tested. Known quantity of protein samples were electrophoresed using Novex® NuPAGE® 12 % Bis-Tris gel, XCell SureLock™ Electrophoresis System and Novex® Sharp Pre-Stained Protein Standard. Resolved proteins were then transferred onto a nitrocellulose membrane with iBlot® 2 Dry Blotting System. The membrane was probed with the relevant primary and secondary Antibody following blocking with 5 % skimmed milk. Chemiluminescent detection was performed using Pierce™ ECL Western Blotting Substrate.




Anti-Caveolin-1 antibody - Caveolae Marker (ab2910) at 2 µg/ml + Rat heart protein extract

**Predicted band size:** 20 kDa




Immunofluorescence analysis of Caveolin 1 was done on 70% confluent log phase A-375 cells. The cells were fixed with 4% paraformaldehyde for 15 minutes, permeabilized with 0.25% Triton™ X-100 for 10 minutes, and blocked with 5% BSA for 1 hour at room temperature. The cells were labeled with ab2910 at 1 µg/mL in 1% BSA and incubated for 3 hours at room temperature and then labeled with Goat anti-Rabbit IgG (H+L) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate at a dilution of 1:2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI. F-actin (Panel c: red) was stained with Alexa Fluor® 555 Rhodamine Phalloidin. Panel d is a merged image showing cytoplasmic localization. Panel e is a no primary antibody control. The images were captured at 60X magnification.



Rat astrocytes stained with fluorescently labeled Caveolin-1 antibody.

Primary antibody is ab2910 at a dilution of 1/500 and the secondary antibody is Texas red labeled anti-rabbit IgG at a dilution of 1/1000.

This image was kindly supplied as part of the review submitted by Donghui Zhu.



Immunocytochemistry/ Immunofluorescence - Anti-Caveolin-1 antibody - Caveolae Marker (ab2910)

This image is courtesy of an anonymous Abreview

ab2910 staining Caveolin-1 - Caveolae Marker in HeLa (Human epithelial cell line from cervix adenocarcinoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence).

Cells were fixed with paraformaldehyde. Samples were incubated with primary antibody (1/200 in PBS + 0.05% Saponin) for 1 hour at 37°C. A Cy3<sup>®</sup>-conjugated Donkey anti-rabbit polyclonal (1/500) was used as the secondary antibody.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors