abcam

Product datasheet

Anti-CD22 (phospho Y762) antibody [EP727Y] ab45135

RabMAb

画像数1

製品の概要

免疫原

製品名 Anti-CD22 (phospho Y762) antibody [EP727Y]

製品の詳細 Rabbit monoclonal [EP727Y] to CD22 (phospho Y762)

由来種 Rabbit

特異性 This antibody will detect CD22 phosphorylation on Tyrosine 762. This antibody is predicted to

detect splice isoform 2 based on sequence homology.

アプリケーション **適用あり:** WB

適用なし: Flow Cyt,ICC/IF or IP

種交差性 交差種: Human

交差が予測される動物種: Goat, Chimpanzee, Monkey, Opossum

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール WB: Raji treated with Pervanadate whole cell lysate

特記事項 Our RabMAb® technology is a patented hybridoma-based technology for making rabbit

monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー pH: 7.20

Preservative: 0.05% Sodium azide

Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue

culture supernatant

精製度 Tissue culture supernatant

ポリ/モノ モノクローナル

クローン名 EP727Y

アイソタイプ lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab45135の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| WB | | 1/5000. Predicted molecular weight: 150 kDa. |

追加情報 Is unsuitable for Flow Cyt,ICC/IF or IP.

ターゲット情報

機能 Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid

tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block

signal transduction through dephosphorylation of signaling molecules.

組織特異性 B-lymphocytes.

配列類似性 Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid binding Ig-like lectin) family.

Contains 6 lg-like C2-type (immunoglobulin-like) domains. Contains 1 lg-like V-type (immunoglobulin-like) domain.

ドメイン Contains 4 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based

inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The

phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.

翻訳後修飾 Phosphorylation of Tyr-762, Tyr-807 and Tyr-822 are involved in binding to SYK, GRB2 and SYK,


respectively. Phosphorylation of Tyr-842 is involved in binding to SYK, PLCG2 and

PIK3R1/PIK3R2.

Phosphorylated on tyrosine residues by LYN.

細胞内局在 Cell membrane.

画像



All lanes : Anti-CD22 (phospho Y762) antibody [EP727Y] (ab45135) at 1/1000 dilution (Unpurified)

Lane 1 : Raji (Human Burkitt's lymphoma B lymphocyte) whole cell lysate

Lane 2: Raji treated with Pervanadate whole cell lysate

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit lgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution

Predicted band size: 150 kDa **Observed band size:** 150 kDa

Please note: All products are "FOR RESEARCH USE ONLY, NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors