abcam

Product datasheet

Anti-CD27 antibody [LG.7F9] - Low endotoxin, Azide free ab171204

画像数1

製品の概要

製品名 Anti-CD27 antibody [LG.7F9] - Low endotoxin, Azide free

製品の詳細 Armenian hamster monoclonal [LG.7F9] to CD27 - Low endotoxin, Azide free

由来種Armenian hamsterアプリケーション適用あり: Flow Cyt種交差性交差種: Mouse

免疫原 The details of the immunogen for this antibody are not available.

ポジティブ・コントロール BALB/c splenocytes.

特記事項 Endotoxin Level: Less than 0.001 ng/μg antibody, as determined by the LAL assay.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー pH: 7.2

Constituent: PBS

Aqueous buffer, no sodium azide.

キャリア・フリー はい

精製度 Protein G purified

ポリ/モノ モノクローナル

クローン名 LG.7F9

1

アプリケーション


The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab171204の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| Flow Cyt | | Use $0.5\mu g$ for 10^{5-8} cells. In a final volume of $100~\mu l$. ab18479 - Armenian Hamster monoclonal lgG, is suitable for use as an isotype control with this antibody. |

ターゲット情報

| 機能 | Receptor for CD70/CD27L. May play a role in survival of activated T-cells. May play a role in apoptosis through association with SNA1. | |
|-------|--|--|
| 組織特異性 | Found in most T-lymphocytes. | |
| 配列類似性 | Contains 3 TNFR-Cys repeats. | |
| 翻訳後修飾 | Phosphorylated and O-glycosylated. | |
| 細胞内局在 | Membrane. | |

画像



Flow cytometric analysis of BALB/c splenocytes with 0.25 μ g Armenian Hamster lgG Isotype Control (open histogram) or 0.25 μ g ab171204 (filled histogram) followed by Anti-Armenian Hamster lgG FITC. Total viable cells were analyzed.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors