

# Anti-CD31 antibody [EPR3094] - BSA and Azide free ab207090

リコンビナント **RabMAb**

★★★★★ **2 Abreviews** **6 References** 画像数 **15**

### 製品の概要

製品名	Anti-CD31 antibody [EPR3094] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EPR3094] to CD31 - BSA and Azide free
由来種	Rabbit
アプリケーション	<b>適用あり:</b> WB, IHC-P, ICC/IF <b>適用なし:</b> Flow Cyt or IP
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: THP-1 and Jurkat cell lysates IHC-P: Human kidney tissue and human muscle tissue ICC/IF: Jurkat cells
特記事項	<p>ab207090 is the carrier-free version of <b>ab76533</b>.</p> <p>This antibody shows no cross-reactivity with rat and mouse samples in WB. However it can give some non specific staining on mouse smooth muscle tissues. Please contact our Scientific support for more information.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <b><a href="#">see here</a></b>.</p>

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

## 製品の特性

### 製品の状態


Liquid

### 保存方法

Shipped at 4°C. Store at +4°C. Do Not Freeze.

### 解離定数 (K<sub>D</sub> 値)

K<sub>D</sub> = 1.79 x 10<sup>-10</sup> M



### バッファー

pH: 7.20

Constituent: PBS

### キャリア・フリー

はい

### 精製度

Protein A purified

### ポリ/モノ

モノクローナル

### クローン名

EPR3094

### アイソタイプ

IgG

## アプリケーション

### The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab207090の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 83 kDa. Treat samples with PNGase F or phosphatase to confirm the specificity of bands if necessary. The observed band size of CD31 may not be the same as predicted MWs in WB due to the different forms and modifications of CD31.
IHC-P	★★★★★ (2)	Use at an assay dependent concentration. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. The ideal fixation time will depend on the size of the tissue block and the type of tissue, but fixation between 18–24h is suitable for most samples. Positive Control: Human tonsil tissue

アプリケーション	Abreviews	特記事項
ICC/IF		<p>Use at an assay dependent concentration.</p> <p>It is recommended to incubate cells with 0.1% Triton-X for 5 min to detect nuclear antigen. Use 0.3M glycine to quench autofluorescence caused by aldehydes.</p> <p>Positive Control: HUVEC cells</p>

#### 追加情報

Is unsuitable for Flow Cyt or IP.

#### ターゲット情報

##### 機能

Induces susceptibility to atherosclerosis (By similarity). Cell adhesion molecule which is required for leukocyte transendothelial migration (TEM) under most inflammatory conditions. Tyr-690 plays a critical role in TEM and is required for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC) and is also essential for the LBRC membrane to be targeted around migrating leukocytes. Prevents phagocyte ingestion of closely apposed viable cells by transmitting 'detachment' signals, and changes function on apoptosis, promoting tethering of dying cells to phagocytes (the encounter of a viable cell with a phagocyte via the homophilic interaction of PECAM1 on both cell surfaces leads to the viable cell's active repulsion from the phagocyte. During apoptosis, the inside-out signaling of PECAM1 is somehow disabled so that the apoptotic cell does not actively reject the phagocyte anymore. The lack of this repulsion signal together with the interaction of the eat-me signals and their respective receptors causes the attachment of the apoptotic cell to the phagocyte, thus triggering the process of engulfment). Isoform Delta15 is unable to protect against apoptosis. Modulates BDKRB2 activation. Regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in human umbilical cord vein cells (HUVEC).

##### 組織特異性

Expressed on platelets and leukocytes and is primarily concentrated at the borders between endothelial cells. Isoform Long predominates in all tissues examined. Isoform Delta12 is detected only in trachea. Isoform Delta14-15 is only detected in lung. Isoform Delta14 is detected in all tissues examined with the strongest expression in heart. Isoform Delta15 is expressed in brain, testis, ovary, cell surface of platelets, human umbilical vein endothelial cells (HUVECs), Jurkat T-cell leukemia, human erythroleukemia (HEL) and U937 histiocytic lymphoma cell lines (at protein level).

##### 配列類似性

Contains 6 Ig-like C2-type (immunoglobulin-like) domains.

##### ドメイン

The Ig-like C2-type domains 2 and 3 contribute to formation of the complex with BDKRB2 and in regulation of its activity.


##### 翻訳後修飾

Phosphorylated on Ser and Tyr residues after cellular activation. In endothelial cells Fyn mediates mechanical-force (stretch or pull) induced tyrosine phosphorylation.

##### 細胞内局在


Membrane. Cell junction. Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells and Cell junction. Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells.

#### 画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody  
[EPR3094] - BSA and Azide free (ab207090)

Immunohistochemical analysis of formalin fixed paraffin embedded human kidney labelling CD31 with ab207090 at a concentration of 1µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with an OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was conducted for 32min with ULTRA cell conditioning solution (CC1 pH8.5). ab207090 anti CD31 antibody was incubated at 37oC for 16min. Sections were counterstained is with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.



Multiplex immunohistochemistry - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76533](#)).

Panel A: merged staining of anti-CD31 (gray; Opal™690), anti-CD3 (green; Opal™520) and anti-CD19 (red; Opal™570) on Formalin/PFA-fixed paraffin-embedded sections of human tonsil. Secondary antibody was Opal Polymer HRP Ms + Rb, nuclear counterstain was DAPI.


Panel B: anti-CD3 stained on T cells with [ab16669](#) at 1/500 dilution

Panel C: anti-CD19 stained on B cells with [ab237772](#) at 1/5000 dilution

Panel D: anti-CD31 stained on endothelial cells and immune cell subsets with ab207090 at 1/500 dilution

The section was incubated in three rounds of staining: in the order of ab207090 and [ab16669](#) for 30 mins, then [ab237772](#) for 10 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76533](#)).

Panel A: merged staining of anti-CD31 (gray; Opal™690), anti-CD3 (green; Opal™520) and anti-CD19 (red; Opal™570) on Formalin/PFA-fixed paraffin-embedded sections of human spleen. Secondary antibody was Opal Polymer HRP Ms + Rb, nuclear counterstain was DAPI.


Panel B: anti-CD3 stained on T cells with [ab16669](#) at 1/500 dilution

Panel C: anti-CD19 stained on B cells with [ab237772](#) at 1/5000 dilution

Panel D: anti-CD31 stained on endothelial cells with ab207090 at 1/500 dilution

The section was incubated in three rounds of staining: in the order of ab207090 and [ab16669](#) for 30 mins, then [ab237772](#) for 10 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.


The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



[ab76533](#) staining CD31 in human kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) [ab97051](#) was used as the secondary antibody at a dilution of 1/500.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76533](#)).





Western blot - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

Anti-CD31 antibody [EPR3094] ([ab76533](#)) at 1/10000 dilution + THP-1 (Human monocytic leukemia cell line) whole cell lysate at 20 µg


### Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 83 kDa

### Blocking and diluting buffer: 5% NFDM /TBST

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76533](#)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

Image from Hofmann NA et al. PLoS One. 2012;7(9):e44468. doi: 10.1371/journal.pone.0044468. Epub 2012 Sep 7. Fig 3.; doi:10.1371/journal.pone.0044468; September 7 2012 PLoS ONE 7(9): e44468.

Immunohistochemical analysis of endothelial colony forming progenitor cell plugs, staining CD31 (green) with [ab76533](#).

Following antigen retrieval and blocking, sections were incubated with primary antibody (1/1000) overnight at 4°C. A Cy5®-conjugated anti-rabbit IgG (2 µg/ml) was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76533](#)).




Immunocytochemistry/ Immunofluorescence - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

Immunocytochemistry analysis of Jurkat (human T cell leukemia T lymphocyte) cells labeling CD31 with **ab76533** at 1/500 (4.9 µg/mL). **ab150077**, AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 (2 µg/mL) was used as the secondary antibody. Cells were fixed with 100% Methanol. DAPI (blue) was used as nuclear counterstain.

Confocal image showing positive staining on Jurkat cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76533**).



Western blot - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

**All lanes** : Anti-CD31 antibody [EPR3094] (**ab76533**) at 1/20000 dilution

**Lane 1** : THP-1 cell lysate

**Lane 2** : Jurkat cell lysate


Lysates/proteins at 10 µg per lane.

#### Secondary

**All lanes** : HRP labelled goat anti-rabbit at 1/2000 dilution

**Predicted band size:** 83 kDa

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76533**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

Clone EPR3094 (ab207090) has been successfully conjugated by Abcam. This image was generated using Anti-CD31 antibody [EPR3094] (Alexa Fluor® 647). Please refer to [ab218582](#) for protocol details.


IHC image of CD31 staining in a section of formalin-fixed paraffin-embedded normal human kidney\*.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) in a Biocare Medical NxGen pressure cooker using retrieval settings of 110°C for 8 minutes. Non-specific protein-protein interactions were then blocked in TBS containing 0.025% (v/v) Triton X-100, 0.3M (w/v) glycine and 1% (w/v) BSA for 1h at room temperature. The section was then incubated overnight at +4°C in TBS containing 0.025% (v/v) Triton X-100 and 1% (w/v) BSA with [ab218582](#) at 1/100 dilution (shown in red) and counterstained using [ab195887](#), Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

For other IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antigen retrieval conditions, antibody concentrations and incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

Immunohistochemical analysis of paraffin embedded human kidney tissue using [ab76533](#) at a 1/250 dilution. Note positive staining of endothelial cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76533](#)).

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)

Immunohistochemical analysis of paraffin embedded human muscle tissue using **ab76533** at a 1/250 dilution. Note positive staining of endothelial cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76533**).

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [EPR3094] - BSA and Azide free (ab207090)  
This image is courtesy of an anonymous Abreview.

**ab76533** staining CD31 in human muscle tissue by

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue was fixed with formaldehyde and a heat mediated antigen retrieval step was performed using citrate buffer pH 6.0. Samples were then blocked with 3% serum for 30 minutes at 20°C followed by incubation with the primary antibody at a 1/200 dilution for 12 hours at 20°C. A Cy3®-conjugated goat anti-rabbit polyclonal was used as secondary antibody at a 1/200 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76533**).



SPR Scanning - Anti-CD31 antibody [EPR3094] -  
BSA and Azide free (ab207090)

Equilibrium dissociation constant ( $K_D$ ) measurement to  
determine antibody affinity to the target antigen.

[Click here to learn more about  \$K\_D\$](#)

This data was developed using the same antibody clone in a  
different buffer formulation containing PBS, BSA, glycerol, and  
sodium azide (**ab76533**).

Tissue Microarray (TMA) data for ab76533			
Normal tissue samples		Malignant tissue samples	
Human cardiac muscle	Human placenta	Clear cell carcinoma of human kidney	Human glioma
Human cerebrum	Human skeletal muscle	Human bladder cancer	Human hepatocellular carcinoma
Human colon	Human skin	Human breast carcinoma	Human lung carcinoma
Human endometrium	Human spleen	Human cervical carcinoma	Human ovarian carcinoma
Human kidney	Human stomach	Human colon carcinoma	Human pancreatic carcinoma
Human liver	Human testis	Human endometrial carcinoma	Human prostatic hyperplasia
Human lung	Human thyroid	Human gastric carcinoma	Human thyroid carcinoma
Human mammary gland	Human tonsil		
Human pancreas			

Immunohistochemistry (Formalin/PFA-fixed paraffin-  
embedded sections) - Anti-CD31 antibody  
[EPR3094] - BSA and Azide free (ab207090)

Tissue Microarrays stained for " Anti-CD31 antibody [EPR3094]"  
using "**ab76533**" in immunohistochemical analysis. This table  
provides a detailed overview of positive (tick mark) and negative  
(cross mark) staining per sample type tested. The sections were  
pre-treated using Heat mediated antigen retrieval using **ab93684**  
(Tris/EDTA buffer, pH 9.0). The sections were incubated with  
**ab76533** at +4°C overnight followed by a ready to use Goat Anti-  
Rabbit IgG H&L (HRP polymer).

Why choose a  
recombinant antibody?

**Research with confidence**  
Consistent and reproducible results

**Long-term and scalable supply**  
Recombinant technology

**Success from the first experiment**  
Confirmed specificity

**Ethical standards compliant**  
Animal-free production

Anti-CD31 antibody [EPR3094] - BSA and Azide  
free (ab207090)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors