

# Anti-CD42d/GP5 antibody [1C2] ab95780

[2 References](#) [画像数 1](#)

### 製品の概要

---

製品名	Anti-CD42d/GP5 antibody [1C2]
製品の詳細	Armenian hamster monoclonal [1C2] to CD42d/GP5
由来種	Armenian hamster
アプリケーション	<b>適用あり:</b> Flow Cyt
種交差性	<b>交差種:</b> Mouse
免疫原	Tissue, cells or virus corresponding to Mouse CD42d/GP5. (C57BL/6 x DBA/2)F1 mouse platelets
ポジティブ・コントロール	Mouse platelets
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide Constituent: 2.68% PBS
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	1C2
アイソタイプ	IgG

アプリケーション


**The Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab95780の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 0.125-0.25µg for 10 <sup>5-8</sup> cells. Staining the cell sample in a final volume of 100 µl is recommended.

### ターゲット情報

<b>機能</b>	The GPIb-V-IX complex functions as the vWF receptor and mediates vWF-dependent platelet adhesion to blood vessels. The adhesion of platelets to injured vascular surfaces in the arterial circulation is a critical initiating event in hemostasis.
<b>組織特異性</b>	Platelets and megakaryocytes.
<b>配列類似性</b>	Contains 13 LRR (leucine-rich) repeats. Contains 1 LRRCT domain. Contains 1 LRRNT domain.
<b>翻訳後修飾</b>	The N-terminus is blocked.
<b>細胞内局在</b>	Membrane.

### 画像



Flow cytometric staining of mouse platelets with Anti-Mouse CD41 FITC and 0.125µg of Armenian Hamster IgG Isotype Control Purified (left) or 0.125µg of ab95780 (right) followed by Anti-Armenian Hamster IgG PE. Total events were used for analysis.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors