# abcam

## Product datasheet

# Anti-CD57 antibody [HNK-1 or Leu-7] - BSA and Azide free ab212403

★★★★★ 1 Abreviews 1 References 画像数 1

#### 製品の概要

製品名 Anti-CD57 antibody [HNK-1 or Leu-7] - BSA and Azide free

製品の詳細 Mouse monoclonal [HNK-1 or Leu-7] to CD57 - BSA and Azide free

由来種 Mouse

 アプリケーション
 適用あり: IHC-P

 種交差性
 交差種: Human

非交差種: Rat

免疫原 Tissue, cells or virus corresponding to Human CD57. Human peripheral blood mononuclear cells.

ポジティブ・コントロール CCRF-CEM cells; Human lymph node or tonsil tissue.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

**バッファー** pH: 7.2

Constituent: 100% PBS

キャリア・フリー はい

精製度 Protein A/G purified

特記事項(精製) ab212403 is purified from Bioreactor Concentrate by Protein A/G.

**ポリ/モノ** モノクローナル **クローン名** HNK-1 or Leu-7

1

lgΜ

軽鎖の種類

kappa

#### アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab212403の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 2 - 4 µg/ml. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. for 30 minutes at RT

#### ターゲット情報

#### 関連性

CD57 is a carbohydrate epitope that contains a sulfoglucuronyl residue in several adhesion molecules of the nervous system. Is is created by Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1 (B3GAT1) The CD57 antigen is present on subsets of T cells and natural killer cells.

#### 画像



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD57 antibody [HNK-1 or Leu-7] - BSA and Azide free (ab212403)

Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human tonsil tissue labeling CD57 with ab212403 at 4µg/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors