

# Anti-CD57 antibody [NK/804] ab220187

**2 References**   [画像数 1](#)

### 製品の概要

製品名	Anti-CD57 antibody [NK/804]
製品の詳細	Mouse monoclonal [NK/804] to CD57
由来種	Mouse
アプリケーション	<b>適用あり:</b> IHC-P
種交差性	<b>交差種:</b> Human <b>非交差種:</b> Rat
免疫原	Recombinant full length protein corresponding to Human CD57. Database link: <a href="#">Q9P2W7</a>
ポジティブ・コントロール	Human tonsil tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.05% Sodium azide Constituents: 99% PBS, 0.05% BSA
特記事項 (精製)	ab220187 is purified from Bioreactor Concentrate.
ポリ/モノ	モノクローナル
クローン名	NK/804
アイソタイプ	IgM
軽鎖の種類	kappa

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、 次のテスト済みアプリケーションにおけるab220187の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 2 - 4 µg/ml. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. for 30 minutes at RT

## ターゲット情報

**関連性** CD57 is a carbohydrate epitope that contains a sulfoglucuronyl residue in several adhesion molecules of the nervous system. Is is created by Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1 (B3GAT1) The CD57 antigen is present on subsets of T cells and natural killer cells.

## 画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-B3GAT1 antibody  
[NK/804] (ab220187)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue labelling B3GAT1 with ab220187 at 4 µg/mL for 30 minutes at room temperature. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at room temperature for 20 minutes.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors