abcam

Product datasheet

Anti-CD59 antibody [EPR22394-242] - BSA and Azide free ab254194

יובעדער RabMAb

画像数 2

製品の概要

Anti-CD59 antibody [EPR22394-242] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EPR22394-242] to CD59 - BSA and Azide free

Rabbit

適用あり: Flow Cyt

適用なし: ICC/IF,IHC-P or WB

交差種: Human

Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

Flow Cyt: Human primary peripheral blood mononuclear cells.

ab254194 is the carrier-free version of ab253239.

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

1

製品名

由来種

アプリケーション

種交差性

免疫原

ポジティブ・コントロール

特記事項

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

バッファー pH: 7.2

Constituent: PBS

キャリア・フリー はい

精製度 Protein A purified

ポリ/モノ モノクローナル

クローン名 EPR22394-242

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab254194の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration.

追加情報

Is unsuitable for ICC/IF,IHC-P or WB.

ターゲット情報

機能

Potent inhibitor of the complement membrane attack complex (MAC) action. Acts by binding to the C8 and/or C9 complements of the assembling MAC, thereby preventing incorporation of the multiple copies of C9 required for complete formation of the osmolytic pore. This inhibitor appears to be species-specific. Involved in signal transduction for T-cell activation complexed to a protein tyrosine kinase.

The soluble form from urine retains its specific complement binding activity, but exhibits greatly reduced ability to inhibit MAC assembly on cell membranes.

関連疾患

Defects in CD59 are the cause of CD59 deficiency (CD59D) [MIM:612300].

配列類似性


Contains 1 UPAR/Ly6 domain.

翻訳後修飾

N- and O-glycosylated. The N-glycosylation mainly consists of a family of biantennary complex-type structures with and without lactosamine extensions and outer arm fucose residues. Also significant amounts of triantennary complexes (22%). Variable sialylation also present in the Asn-43 oligosaccharide. The predominant O-glycans are mono-sialylated forms of the disaccharide, Gal-beta-1,3GalNAc, and their sites of attachment are probably on Thr-76 and Thr-77. The GPl-anchor of soluble urinary CD59 has no inositol-associated phospholipid, but is composed of seven different GPl-anchor variants of one or more monosaccharide units. Major variants contain sialic acid, mannose and glucosamine Sialic acid linked to an N-acetylhexosamine-galactose arm is present in two variants.

Glycated. Glycation is found in diabetic subjects, but only at minimal levels in nondiabetic subjects. Glycated CD59 lacks MAC-inhibitory function and confers to vascular complications of diabetes.

画像



Flow Cytometry - Anti-CD59 antibody [EPR22394-242] - BSA and Azide free (ab254194) Flow cytometric analysis of U-937 (human histiocytic lymphoma monocyte, Left panel) / Human primary peripheral blood mononuclear cell (PBMC, Right panel) labeling CD59 with ab253239 at 1/400 dilution (red) compared with a Rabbit monoclonal lgG (ab172730) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat anti rabbit lgG (Alexa Fluor® 488, ab150077), at 1/5000 dilution was used as the secondary antibody.

Negative control: U-937 (PMID:16493049). Gated on viable cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab253239).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

• We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors