abcam

Product datasheet

Anti-CD62P antibody [EPR22850-206] - BSA and Azide free ab259996

יובעדער RabMAb

画像数 2

製品の概要

製品名 Anti-CD62P antibody [EPR22850-206] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EPR22850-206] to CD62P - BSA and Azide free

由来種 Rabbit

アプリケーション 適用あり: Flow Cyt

適用なし: ICC/IF,IHC-P,IP or WB

種交差性 交差種: Human

免疫原 Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール Flow Cyt: HEK-293 cells.

特記事項 ab259996 is the carrier-free version of ab255821.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

1

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

バッファー pH: 7.2

Constituent: PBS

キャリア・フリー はい

精製度 Protein A purified

ポリ/モノ モノクローナル

クローン名 EPR22850-206

アイソタイプ lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab259996の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration.

追加情報 Is unsuitable for ICC/IF,IHC-P,IP or WB.

ターゲット情報

機能 Ca(2+)-dependent receptor for myeloid cells that binds to carbohydrates on neutrophils and

monocytes. Mediates the interaction of activated endothelial cells or platelets with leukocytes. The ligand recognized is sialyl-Lewis X. Mediates rapid rolling of leukocyte rolling over vascular

surfaces during the initial steps in inflammation through interaction with PSGL1.

組織特異性 Stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. Upon cell

activation by agonists, P-selectin is transported rapidly to the cell surface.

関連疾患 Defects in SELP may be a cause of susceptibility to ischemic stroke (ISCHSTR) [MIM:601367];

also known as cerebrovascular accident or cerebral infarction. A stroke is an acute neurologic event leading to death of neural tissue of the brain and resulting in loss of motor, sensory and/or cognitive function. Ischemic strokes, resulting from vascular occlusion, is considered to be a highly

complex disease consisting of a group of heterogeneous disorders with multiple genetic and

environmental risk factors.


配列類似性 Belongs to the selectin/LECAM family.

Contains 1 C-type lectin domain.
Contains 1 EGF-like domain.

Contains 9 Sushi (CCP/SCR) domains.

細胞内局在 Membrane.

画像




Flow Cytometry - Anti-CD62P antibody [EPR22850-206] - BSA and Azide free (ab259996)

Flow cytometric analysis of HEK-293 (human embryonic kidney epithelial cell, Left) / HEL (human Erythroleukemia erythroblast, Right) cells labelling CD62P with **ab255821** at 1/50 dilution (Red) compared with a Rabbit monoclonal lgG (**ab172730**) / Black isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit lgG (Alexa Fluor[®] 488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.

Negative control: HEK-293 (PMID: 7693674).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab255821</u>).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors