

Anti-CD8 alpha antibody [BLR044F] - BSA free ab272093

リコンビナント

画像数 5

製品の概要

製品名	Anti-CD8 alpha antibody [BLR044F] - BSA free
製品の詳細	Rabbit monoclonal [BLR044F] to CD8 alpha - BSA free
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human
免疫原	Synthetic peptide within Human CD8 alpha aa 185-235 (C terminal). The exact sequence is proprietary. NP_001759.3 and Gene ID 925. Database link: P01732
特記事項	ab272093 is the BSA-free version of ab243900 . This product is sold under License from Bethyl Laboratories, Inc.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 8.20 Preservative: 0.09% Sodium azide Constituent: 98% Borate buffered saline
特記事項(精製)	Recombinant antibody was purified from cell culture supernatant.
ポリ/モノ	モノクローナル
クローン名	BLR044F
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab272093の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報

機能	Identifies cytotoxic/suppressor T-cells that interact with MHC class I bearing targets. CD8 is thought to play a role in the process of T-cell mediated killing. CD8 alpha chains binds to class I MHC molecules alpha-3 domains.
関連疾患	Defects in CD8A are a cause of familial CD8 deficiency (CD8 deficiency) [MIM:608957]. Familial CD8 deficiency is a novel autosomal recessive immunologic defect characterized by absence of CD8+ cells, leading to recurrent bacterial infections.
配列類似性	Contains 1 Ig-like V-type (immunoglobulin-like) domain.
翻訳後修飾	All of the five most carboxyl-terminal cysteines form inter-chain disulfide bonds in dimers and higher multimers, while the four N-terminal cysteines do not.
細胞内局在	Secreted and Cell membrane.

画像



All lanes : Anti-CD8 alpha antibody [BLR044F] ([ab243900](#)) at 1/1000 dilution

Lane 1 : CD4+ T cell whole cell lysate

Lane 2 : CD8+ T cell whole cell lysate

Lane 3 : Jurkat whole cell lysate

Lane 4 : MOLT-4 whole cell lysate

Lane 5 : RPMI 8226 whole cell lysate

Lane 6 : MJ whole cell lysate

Lane 7 : C5/MJ whole cell lysate

Lane 8 : KG-1 whole cell lysate

Lane 9 : SR whole cell lysate

Lane 10 : K562 whole cell lysate

Lysates/proteins at 50 µg per lane.


Secondary

All lanes : HRP-conjugated goat anti-rabbit IgG

Detection: Chemiluminescence.

Lysates prepared using NETN lysis buffer.


This data was developed using the same antibody clone in a different buffer formulation containing Borate buffered saline, BSA, glycerol and sodium azide ([ab243900](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD8 alpha antibody [BLR044F] - BSA free ([ab272093](#))

Formalin-fixed, paraffin-embedded human tonsil tissue stained for CD8 alpha using [ab243900](#) at 1/250 dilution in immunohistochemical analysis. A HRP-conjugated goat anti-rabbit antibody was used as the secondary. DAB staining.


This data was developed using the same antibody clone in a different buffer formulation containing Borate buffered saline, BSA, glycerol and sodium azide ([ab243900](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD8 alpha antibody [BLR044F] - BSA free ([ab272093](#))

This data was developed using the same antibody clone in a different buffer formulation containing Borate buffered saline, BSA, glycerol and sodium azide ([ab243900](#)).

Formalin-fixed, paraffin-embedded human melanoma tissue stained for CD8 alpha (red) using [ab243900](#) at 1/100 dilution in immunohistochemical analysis. A conjugated goat anti-rabbit antibody was used as the secondary. Counterstain: DAPI (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD8 alpha antibody [BLR044F] - BSA free (ab272093)

This data was developed using the same antibody clone in a different buffer formulation containing Borate buffered saline, BSA, glycerol and sodium azide (**ab243900**).

Formalin-fixed, paraffin-embedded human breast carcinoma tissue stained for CD8 alpha (red) using **ab243900** at 1/250 dilution in immunohistochemical analysis. A conjugated goat anti-rabbit antibody was used as the secondary. Counterstain: DAPI (blue).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-CD8 alpha antibody [BLR044F] - BSA free (ab272093)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors