

Anti-CD8 alpha antibody [CAL38] ab237723

リコンビナント RabMAb

5 References 画像数 5

製品の概要

製品名	Anti-CD8 alpha antibody [CAL38]
製品の詳細	Rabbit monoclonal [CAL38] to CD8 alpha
由来種	Rabbit
アプリケーション	適用あり: IHC-Fr, IP, IHC-P 適用なし: Flow Cyt
種交差性	交差種: Mouse
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Mouse spleen tissue. IHC-Fr: Mouse Thymus tissue. IP: Mouse thymus tissue lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.05% Sodium azide Constituent: PBS
精製度	Protein A purified
特記事項(精製)	Purity >99%.
ポリ/モノ	モノクローナル
クローン名	CAL38
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab237723の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		1/250. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)
IP		1/20.
IHC-P		1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

追加情報

Is unsuitable for Flow Cyt.

ターゲット情報

機能

Identifies cytotoxic/suppressor T-cells that interact with MHC class I bearing targets. CD8 is thought to play a role in the process of T-cell mediated killing. CD8 alpha chains binds to class I MHC molecules alpha-3 domains.

関連疾患

Defects in CD8A are a cause of familial CD8 deficiency (CD8 deficiency) [MIM:608957]. Familial CD8 deficiency is a novel autosomal recessive immunologic defect characterized by absence of CD8+ cells, leading to recurrent bacterial infections.

配列類似性

Contains 1 Ig-like V-type (immunoglobulin-like) domain.


翻訳後修飾

All of the five most carboxyl-terminal cysteines form inter-chain disulfide bonds in dimers and higher multimers, while the four N-terminal cysteines do not.

細胞内局在

Secreted and Cell membrane.


画像



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling CD8 alpha with ab237723 at 1/500 dilution, followed by the ready to use Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).

Positive staining on the mouse spleen is shown. The section was incubated with ab237723 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND[®] RX instrument.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is the ready to use Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).



Immunohistochemistry (Frozen sections) - Anti-CD8 alpha antibody [CAL38] (ab237723)

Immunohistochemical analysis of 4% paraformaldehyde-fixed, 0.2% Triton X-100 permeabilized frozen mouse thymus tissue labeling CD8 alpha with ab237723 at 1/250 dilution, followed by a AlexaFluor®488 Goat anti-Rabbit secondary antibody (**ab150077**) at 1/1000 dilution. Positive staining on the membrane of T cells in mouse thymus (PMID: 29042640) is observed. Counterstained with DAPI (Blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a AlexaFluor®488 Goat anti-Rabbit secondary antibody (**ab150077**) at 1/1000 dilution.



Immunoprecipitation - Anti-CD8 alpha antibody [CAL38] (ab237723)

CD8 alpha was immunoprecipitated from 0.35 mg of Mouse thymus tissue lysate with ab237723 at 1/20 dilution. Western blot was performed on the immunoprecipitate using ab237723 at 1/1000 dilution. VeriBlot for IP secondary antibody (HRP) (**ab131366**) was used as the secondary antibody at 1/5000 dilution.


Lane 1: Mouse thymus tissue lysate 10µg (input)

Lane 2: ab237723 IP in mouse thymus tissue lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab237723 in Mouse thymus lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 30 seconds.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD8 alpha antibody [CAL38] (ab237723)

Formalin-fixed, paraffin-embedded mouse spleen tissue stained for CD8 alpha with ab237723 at 0.125 µg/ml in immunohistochemical analysis.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-CD8 alpha antibody [CAL38] (ab237723)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors