


Anti-CDKN2A/p16INK4a antibody - N-terminal ab189034

★★★★☆ **5 Abreviews** **58 References** 画像数 4

製品の概要

製品名	Anti-CDKN2A/p16INK4a antibody - N-terminal
製品の詳細	Rabbit polyclonal to CDKN2A/p16INK4a - N-terminal
由来種	Rabbit
特異性	Expression levels of the CDKN2A/p16INK4a protein may vary with sample type. It's barely expressed in normal tissue, and mostly expressed in some tumour tissues, such as cervical cancer, breast cancer and so on. Moreover, only expressed in some cell lines. Please see images for recommended positive controls.
アプリケーション	適用あり: ICC/IF, WB, IHC-P
種交差性	交差種: Mouse, Human 交差が予測される動物種: Rat 
免疫原	Synthetic peptide corresponding to Human CDKN2A/p16INK4a (N terminal). 19 amino acid peptide. Database link: P42771
ポジティブ・コントロール	Human colon tissues. Human colon cells. Mouse colon tissue lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.4 Preservative: 0.02% Sodium azide Constituent: 99% PBS
精製度	Immunogen affinity purified

ポリモノ

ポリクローナル

アイソタイプ

IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab189034の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 20 µg/ml.
WB	★★★★★ (2)	Use a concentration of 1 - 2 µg/ml.
IHC-P	★★★★★ (2)	Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報


細胞内局在

Cytoplasmic and Nuclear

製品の状態

There are 4 isoforms produced by alternative splicing. Isoform 1 also known as: p16INK4a; Isoform 3 also known as: p12; Isoform 4 also known as: p14ARF; p19ARF; ARF.

画像



All lanes : Left: ab189034 at 2 ug/mL
Right: **ab270058** at 1000 dilution

Lane 1 : HeLa cell lysate
Lane 2 : HEK-293 cell lysate
Lane 3 : Saos-2 cell lysate
Lane 4 : MCF7 cell lysate


Lysates/proteins at 20 µg per lane.

Developed using the ECL technique.

Performed under reducing conditions.


Observed band size: 17 kDa

Western blot: Anti-CDKN2A antibody (ab189034) staining at 2 ug/ml, shown in green; Mouse anti-GAPDH antibody [6C5] (**ab8245**) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab189034 was shown to bind specifically to CDKN2A however we recommend using **ab270058** for more specific binding in Western blot. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3% milk in TBS-0.1% Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CDKN2A/p16INK4a antibody - N-terminal (ab189034)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue sections labeling CDKN2A/p16INK4a with ab189034 at 5 µg/ml. Heat mediated antigen retrieval was performed.



Immunocytochemistry/ Immunofluorescence - Anti-CDKN2A/p16INK4a antibody - N-terminal (ab189034)

Immunofluorescent analysis of human colon cells labeling CDKN2A using ab189034 at 20 µg/ml.



Immunohistochemical analysis of formalin fixed, paraffin embedded human colon tissue labeling CDKN2A using ab189034 at 5 µg/ml.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CDKN2A/p16INK4a antibody - N-terminal (ab189034)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors