

Anti-Chromogranin A antibody [LK2H10] - BSA and Azide free ab212290

画像数 2

製品の概要

製品名	Anti-Chromogranin A antibody [LK2H10] - BSA and Azide free
製品の詳細	Mouse monoclonal [LK2H10] to Chromogranin A - BSA and Azide free
由来種	Mouse
アプリケーション	適用あり: IHC-P
種交差性	交差種: Human 非交差種: Sheep, Rabbit, Guinea pig
免疫原	Tissue, cells or virus corresponding to Human Chromogranin A. (Human pheochromocytoma).
ポジティブ・コントロール	Human adrenal gland and small cell lung carcinoma tissues; PC12 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Constituent: 100% PBS
キャリア・フリー	はい
精製度	Protein A/G purified
特記事項(精製)	ab212290 was purified from Bioreactor Concentrate by Protein A/G.
ポリ/モノ	モノクローナル
クローン名	LK2H10

アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab212290の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 0.25 - 0.5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. (Primary incubation for 30 minutes at RT).

ターゲット情報


機能	Pancreastatin strongly inhibits glucose induced insulin release from the pancreas.
配列類似性	Belongs to the chromogranin/secretogranin protein family.
翻訳後修飾	Sulfated on tyrosine residues and/or contains sulfated glycans. O-glycosylated with core 1 or possibly core 8 glycans.
細胞内局在	Secreted. Neuroendocrine and endocrine secretory granules.
製品の状態	According to the Swiss-Prot database, this protein has many different forms which correspond to various molecular weights; thus the possible variation from the predicted molecular weight of 51kDa.

画像



Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human adrenal gland tissue labeling Chromogranin A with ab212290 at 0.5 µg/ml.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Chromogranin A antibody [LK2H10] - BSA and Azide free (ab212290)



Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human small cell lung carcinoma tissue labeling Chromogranin A with ab212290 at 0.5 µg/ml.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Chromogranin A antibody [LK2H10] - BSA and Azide free (ab212290)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors