

Anti-Collagen III antibody [FH-7A] ab6310

★★★★★ [14 Abreviews](#) [162 References](#) [画像数 3](#)

製品の概要

製品名	Anti-Collagen III antibody [FH-7A]
製品の詳細	Mouse monoclonal [FH-7A] to Collagen III
由来種	Mouse
特異性	ab6310 specifically recognizes collagen type III from human and rat origin. It does not recognize collagen types I, II, IV, V, VI and X.
アプリケーション	適用あり: IHC-Fr, IHC-P, WB, ELISA, Dot blot, Indirect ELISA, ICC/IF
種交差性	交差種: Rat, Human
免疫原	Full length native protein (purified) (Human).
ポジティブ・コントロール	IHC-P: Rat skin sections. IHC-Fr: Rat skin sections.
特記事項	Type III collagen, [α1(III)] ₃ , is an approx. 300 kDa molecule, found predominantly in skin, blood vessels, liver, placenta, tongue, and thymus. Collagen type III forms cofibrils with type I and/or V collagens in a number of tissues of mesenchymal origin, such as skin, tendon, ligaments, and bone. This collagen type is involved, directly or indirectly in several genetic diseases, including Ehlers-Danlos type IV disease.

This product was changed from ascites to tissue culture supernatant on 17 May 2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.

バッファー	Preservative: 0.097% Sodium azide Constituent: Whole serum
精製度	Tissue culture supernatant
ポリ/モノ	モノクローナル
クローン名	FH-7A
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab6310の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		Use at an assay dependent concentration.
IHC-P	★★★★★ (10)	Use at an assay dependent concentration.
WB	★★★★★ (1)	Use at an assay dependent concentration.
ELISA	★★★★☆ (2)	Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
Indirect ELISA		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration. PubMed: 25136258

ターゲット情報

機能	Collagen type III occurs in most soft connective tissues along with type I collagen.
関連疾患	<p>Defects in COL3A1 are a cause of Ehlers-Danlos syndrome type 3 (EDS3) [MIM:130020]; also known as benign hypermobility syndrome. EDS is a connective tissue disorder characterized by hyperextensible skin, atrophic cutaneous scars due to tissue fragility and joint hyperlaxity. EDS3 is a form of Ehlers-Danlos syndrome characterized by marked joint hyperextensibility without skeletal deformity.</p> <p>Defects in COL3A1 are the cause of Ehlers-Danlos syndrome type 4 (EDS4) [MIM:130050]. EDS is a connective tissue disorder characterized by hyperextensible skin, atrophic cutaneous scars due to tissue fragility and joint hyperlaxity. EDS4 is the most severe form of the disease. It is characterized by the joint and dermal manifestations as in other forms of the syndrome, characteristic facial features (acrogeria) in most patients, and by proneness to spontaneous rupture of bowel and large arteries. The vascular complications may affect all anatomical areas.</p> <p>Defects in COL3A1 are a cause of susceptibility to aortic aneurysm abdominal (AAA) [MIM:100070]. AAA is a common multifactorial disorder characterized by permanent dilation of the abdominal aorta, usually due to degenerative changes in the aortic wall. Histologically, AAA is characterized by signs of chronic inflammation, destructive remodeling of the extracellular matrix, and depletion of vascular smooth muscle cells.</p>

配列類似性

Belongs to the fibrillar collagen family.
Contains 1 fibrillar collagen NC1 domain.
Contains 1 VWFC domain.


翻訳後修飾

Proline residues at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.
O-linked glycan consists of a Glc-Gal disaccharide bound to the oxygen atom of a post-translationally added hydroxyl group.

細胞内局在

Secreted > extracellular space > extracellular matrix.


画像



Staining of formalin-fixed, paraffin-embedded rat skin with 1:4,000 ab6310 using biotin/ExtrAvidin®-Peroxidase.

This image was generated using the ascites version of the product.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen III antibody [FH-7A] (ab6310)




ab6310 at 1/600 dilution staining preovulatory follicle and whole ovary tissue sections by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Antigens were retrieved by boiling with an antigen unmasking solution for 20 min at 120°C in an autoclave and then cooled down in water for 5 minutes. The tissue sections were formaldehyde fixed and incubated with the antibody for 1 hour. An alkaline phosphatase conjugated antibody was used as the secondary. The image shows a section of whole preovulatory follicle. Staining for collagen type III is seen in the theca interna cell layer. No staining in the granulosa cells.

This image was generated using the ascites version of the product.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen III antibody [FH-7A] (ab6310)

This image is courtesy of an Abreview submitted by Birgitta Weijdegard



Immunohistochemistry (Frozen sections) - Anti-Collagen III antibody [FH-7A] (ab6310)

Staining of frozen rat skin sections with 1:8,000 ab6310 using biotin/ExtrAvidin®-Peroxidase.

This image was generated using the ascites version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors