

Anti-Collagen XVII antibody [EPR18614] ab184996

リコンビナント **RabMAb**

★★★★★ **4 Abreviews** **25 References** 画像数 12

製品の概要

製品名	Anti-Collagen XVII antibody [EPR18614]
製品の詳細	Rabbit monoclonal [EPR18614] to Collagen XVII
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF, IP
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: A431 whole cell lysate, human, mouse and rat skin lysates. IHC-P: Human esophagus, mouse skin and rat skin tissues. ICC/IF: HaCaT and A431 cells. IP: A431 whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol, 0.05% BSA</p>
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR18614
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab184996の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	1/1000. Detects a band of approximately 180, 130 kDa (predicted molecular weight: 150 kDa).
IHC-P		1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF	★★★★★ (1)	1/100.
IP	★★★★★ (1)	1/50.

ターゲット情報

関連性 Unlike most collagens, collagen XVII is a transmembrane protein. Collagen XVII is a structural component of hemidesmosomes, multiprotein complexes at the dermal epidermal basement membrane zone that mediate adhesion of keratinocytes to the underlying membrane. Mutations in the gene coding for collagen XVII are associated with both generalized atrophic benign and junctional epidermolysis bullosa. Two homotrimeric forms of type XVII collagen exist. The full length form is the transmembrane protein. A soluble form, referred to as either ectodomain or LAD 1, is generated by proteolytic processing of the full length form. Two transcript variants, one resulting from alternative splicing in the 3' UTR, have been identified for this gene.

細胞内局在 Cell junction, hemidesmosome. Membrane; Single-pass type II membrane protein.
Note=Localized along the plasma membrane of the hemidesmosome. 120 kDa linear IgA disease antigen and 97 kDa linear IgA disease antigen: Secreted, extracellular space, extracellular matrix, basement membrane.

画像



All lanes : Anti-Collagen XVII antibody [EPR18614] (ab184996) at 1/5000 dilution

Lane 1 : Human skin lysate

Lane 2 : Human fetal liver lysate

Lane 3 : Human fetal lung lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/10000 dilution


Predicted band size: 150 kDa

Observed band size: 130,180 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDm/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 15161638).



Anti-Collagen XVII antibody [EPR18614] (ab184996) at 1/5000 dilution + A431 (Human epidermoid carcinoma) whole cell lysate at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution


Predicted band size: 150 kDa

Observed band size: 130,180 kDa

Exposure time: 1 minute

Blocking/Dilution buffer: 5% NFDm/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 15161638).



Western blot - Anti-Collagen XVII antibody
[EPR18614] (ab184996)

Anti-Collagen XVII antibody [EPR18614] (ab184996) at 1/2000 dilution + Mouse skin lysate at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution


Predicted band size: 150 kDa

Observed band size: 130,180 kDa

Exposure time: 2 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 15161638).



Western blot - Anti-Collagen XVII antibody
[EPR18614] (ab184996)

Anti-Collagen XVII antibody [EPR18614] (ab184996) at 1/1000 dilution + Rat skin lysate at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution


Predicted band size: 150 kDa

Observed band size: 130,180 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 15161638).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen XVII antibody [EPR18614] (ab184996)

Immunohistochemical analysis of paraffin-embedded Human esophagus tissue labeling Collagen XVII with ab184996 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution. Cytoplasmic staining on squamous epithelial cells of Human esophagus is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

J Biol Chem. 1991. 266, 24064-24069.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen XVII antibody [EPR18614] (ab184996)

Immunohistochemical analysis of paraffin-embedded Human pancreas tissue labeling Collagen XVII with ab184996 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution. Negative on Human pancreas. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen XVII antibody [EPR18614] (ab184996)

Immunohistochemical analysis of paraffin-embedded mouse skin tissue labeling Collagen XVII with ab184996 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution. Cytoplasmic and membrane staining on squamous epithelial cells of mouse skin is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen XVII antibody [EPR18614] (ab184996)

Immunohistochemical analysis of paraffin-embedded rat skin tissue labeling Collagen XVII with ab184996 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution. Cytoplasmic staining on epithelial cells of rat skin is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HaCaT (Human keratinocyte cells) cells labeling Collagen XVII with ab184996 at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic and membranous staining on HaCaT cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with [ab7291](#) (anti-Tubulin mouse mAb) at 1/1000 dilution and [ab150120](#) (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: ab184996 at 1/100 dilution followed by [ab150120](#)

(AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.

-ve control 2: [ab7291](#) (anti-Tubulin mouse mAb) at 1/1000 dilution followed by [ab150077](#) (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized A431 (Human epidermoid carcinoma) cells labeling Collagen XVII with ab184996 at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic and membranous staining on A431 cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with [ab7291](#) (anti-Tubulin mouse mAb) at 1/1000 dilution and [ab150120](#) (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: ab184996 at 1/100 dilution followed by [ab150120](#)

(AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.

-ve control 2: [ab7291](#) (anti-Tubulin mouse mAb) at 1/1000 dilution followed by [ab150077](#) (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Immunoprecipitation - Anti-Collagen XVII antibody
[EPR18614] (ab184996)

Collagen XVII was immunoprecipitated from 1mg of A431 (Human epidermoid carcinoma) whole cell lysate with ab184996 at 1/50 dilution. Western blot was performed from the immunoprecipitate using ab184996 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: A431 whole cell lysate 10ug (Input).

Lane 2: ab184996 IP in A431 whole cell lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab184996 in A431 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Collagen XVII antibody [EPR18614] (ab184996)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors