

Anti-CXXC5 antibody ab106533

2 References [画像数 3](#)

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-CXXC5 antibody |
| 製品の詳細 | Rabbit polyclonal to CXXC5 |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: ICC/IF, IHC-P, WB |
| 種交差性 | 交差種: Human |
| 免疫原 | Synthetic peptide, 15 aa from near the amino terminus of human CXXC5 (NP_057547). |
| ポジティブ・コントロール | Human brain tissue lysate. |
| 特記事項 | <p>Predicted to not cross-react with other CXXC protein family members.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|---------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at 4°C (stable for up to 12 months). |
| バッファー | <p>pH: 7.2</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: PBS</p> |
| 精製度 | Immunogen affinity purified |
| 一次抗体 備考 | Predicted to not cross-react with other CXXC protein family members. |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab106533の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| ICC/IF | | Use a concentration of 20 µg/ml. |
| IHC-P | | Use a concentration of 2.5 µg/ml. |
| WB | | Use a concentration of 1 - 2 µg/ml. Predicted molecular weight: 33 kDa. Isoform 2 predicted molecular weight 24 kDa |

ターゲット情報


| | |
|-------|--|
| 機能 | May indirectly participate in activation of the NF-kappa-B and MAPK pathways. Acts as a mediator of BMP4-mediated modulation of canonical Wnt signaling activity in neural stem cells. |
| 配列類似性 | Contains 1 CXXC-type zinc finger. |
| 細胞内局在 | Nucleus. Cytoplasm. Colocalizes with DVL1 in large bodies localized just outside the nuclear membrane. |
| 製品の状態 | There are 2 isoforms produced by alternative splicing. |

画像



Anti-CXXC5 antibody (ab106533) at 1 µg/ml + human brain tissue lysate at 15 µg/ml

Predicted band size: 33 kDa



Immunohistochemistry of CXXC5 in human brain tissue with ab106533 at 2.5 µg/ml.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CXXC5 antibody (ab106533)

Immunocytochemistry/ Immunofluorescence - Anti-CXXC5 antibody (ab106533)

Immunofluorescence of CXXC5 in Human Brain cells using ab106533 at 20 ug/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors