

Anti-Cytokeratin 1 antibody [EPR17744] ab185628

リコンビナント **RabMAb**

11 References **画像数 8**

製品の概要

製品名	Anti-Cytokeratin 1 antibody [EPR17744]
製品の詳細	Rabbit monoclonal [EPR17744] to Cytokeratin 1
由来種	Rabbit
特異性	Recombinant mouse cytokeratin 5 could be detected at very high antibody concentrations in WB.
アプリケーション	適用あり: WB, IHC-P, IHC-Fr
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Mouse, rat and human skin lysates. IHC-P: Mouse and rat skin tissues. IHC-Fr: Mouse skin tissue.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR17744

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab185628の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/20000. Detects a band of approximately 66 kDa (predicted molecular weight: 66 kDa).
IHC-P		1/600. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IHC-Fr		1/200.

ターゲット情報

機能	May regulate the activity of kinases such as PKC and SRC via binding to integrin beta-1 (ITB1) and the receptor of activated protein kinase C (RACK1/GNB2L1).
組織特異性	The source of this protein is neonatal foreskin. The 67-kDa type II keratins are expressed in terminally differentiating epidermis.
関連疾患	<p>Defects in KRT1 are a cause of bullous congenital ichthyosiform erythroderma (BCIE) [MIM:113800]; also known as epidermolytic hyperkeratosis (EHK) or bullous erythroderma ichthyosiformis congenita of Brocq. BCIE is an autosomal dominant skin disorder characterized by widespread blistering and an ichthyotic erythroderma at birth that persist into adulthood. Histologically there is a diffuse epidermolytic degeneration in the lower spinous layer of the epidermis. Within a few weeks from birth, erythroderma and blister formation diminish and hyperkeratoses develop.</p> <p>Defects in KRT1 are the cause of ichthyosis hystrix Curth-Macklin type (IHCM) [MIM:146590]. IHCM is a genodermatosis with severe verrucous hyperkeratosis. Affected individuals manifest congenital verrucous black scale on the scalp, neck, and limbs with truncal erythema, palmoplantar keratoderma and keratoses on the lips, ears, nipples and buttocks.</p> <p>Defects in KRT1 are a cause of palmoplantar keratoderma non-epidermolytic (NEPPK) [MIM:600962]. NEPPK is a dermatological disorder characterized by focal palmoplantar keratoderma with oral, genital, and follicular lesions.</p> <p>Defects in KRT1 are a cause of ichthyosis annular epidermolytic (AEI) [MIM:607602]; also known as cyclic ichthyosis with epidermolytic hyperkeratosis. AEI is a skin disorder resembling bullous congenital ichthyosiform erythroderma. Affected individuals present with bullous ichthyosis in early childhood and hyperkeratotic lichenified plaques in the flexural areas and extensor surfaces at later ages. The feature that distinguishes AEI from BCIE is dramatic episodes of flares of annular polycyclic plaques with scale, which coalesce to involve most of the body surface and can persist for several weeks or even months.</p> <p>Defects in KRT1 are the cause of palmoplantar keratoderma striate type 3 (SPPK3) [MIM:607654]; also known as keratosis palmoplantaris striata III. SPPK3 is a dermatological disorder affecting palm and sole skin where stratum corneum and epidermal layers are thickened. There is no involvement of non-palmoplantar skin, and both hair and nails are normal.</p>

配列類似性

Belongs to the intermediate filament family.


翻訳後修飾

Undergoes deimination of some arginine residues (citrullination).

細胞内局在

Cell membrane. Located on plasma membrane of neuroblastoma NMB7 cells.

画像




Immunohistochemical analysis of paraffin-embedded mouse skin tissue labeling Cytokeratin 1 with ab185628 at 1/600 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasm staining on mouse skin is observed (PMID: 12010363).

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemical analysis of paraffin-embedded mouse stomach tissue labeling Cytokeratin 1 with ab185628 at 1/600 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Negative Control: No staining on mouse stomach.

Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 1 antibody [EPR17744] (ab185628)


Immunohistochemical analysis of paraffin-embedded rat skin tissue labeling Cytokeratin 1 with ab185628 at 1/600 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasm staining on rat skin is observed (PMID: 12010363).

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 1 antibody [EPR17744] (ab185628)

Immunohistochemical analysis of paraffin-embedded rat kidney tissue labeling Cytokeratin 1 with ab185628 at 1/600 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Negative Control: No staining on rat kidney.

Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Frozen sections) - Anti-Cytokeratin 1 antibody [EPR17744] (ab185628)

Immunohistochemical analysis of 4% paraformaldehyde-fixed, 0.2% Triton X-100 permeabilized frozen mouse skin tissue section labeling Cytokeratin 1 with ab185628 at 1/200 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green).

The result showed cytoplasmic staining on mouse skin.

The nuclear counter stain is DAPI (blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) at 1/1000 dilution.



Western blot - Anti-Cytokeratin 1 antibody [EPR17744] (ab185628)

All lanes : Anti-Cytokeratin 1 antibody [EPR17744] (ab185628) at 1/20000 dilution

Lane 1 : Mouse skin lysate

Lane 2 : Mouse brain lysate

Lane 3 : Mouse spleen lysate

Lane 4 : Mouse heart lysate

Lane 5 : Rat skin lysate

Lane 6 : Rat brain lysate

Lane 7 : Rat spleen lysate

Lane 8 : C6 (Rat glial tumor cell line) whole cell lysate

Lane 9 : RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 66 kDa

Observed band size: 66 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.

Exposure time: Lane 1-4: 30 seconds; Lane 5-9: 3 minutes.

Cytokeratin 1 is expressed in terminally differentiating epidermis (PMID: 2580302).



All lanes : Anti-Cytokeratin 1 antibody [EPR17744] (ab185628) at 1/20000 dilution

Lane 1 : Human skin lysate

Lane 2 : Human fetal brain lysate

Lane 3 : Human fetal heart lysate

Lane 4 : Human fetal kidney lysate

Lane 5 : Human spleen lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 66 kDa

Observed band size: 66 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDm/TBST.

Cytokeratin 1 is expressed in terminally differentiating epidermis (PMID: 2580302).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Cytokeratin 1 antibody [EPR17744] (ab185628)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors