

Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free ab255694

リコンビナント RabMAb

画像数 9

製品の概要

製品名	Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EPR22592-18] to DLL3 - BSA and Azide free
由来種	Rabbit
特異性	The mouse recommendation is based on the IHC-P results. We do not guarantee WB for mouse.
アプリケーション	適用あり: Flow Cyt (Intra), WB, IHC-P, ICC/IF, IP
種交差性	交差種: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HEK-293T transfected with DLL3 (WT) expression vector containing a myc-His-tag®, whole cell lysate. IHC-P: Human small cell lung cancer and glioma tissue; Mouse embryonic brain of day 14.5 tissue. ICC/IF: HEK-293T cells. Flow: HEK-293T cells. IP: HEK-293T transfected with DLL3 expression construct containing a myc-His-tag® whole cell lysate.
特記事項	<p>ab255694 is the carrier-free version of ab229902.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.2 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR22592-18
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab255694の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 75 kDa (predicted molecular weight: 65 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.

ターゲット情報

機能	Inhibits primary neurogenesis. May be required to divert neurons along a specific differentiation pathway. Plays a role in the formation of somite boundaries during segmentation of the paraxial mesoderm.
関連疾患	Spondylocostal dysostosis 1
配列類似性	Contains 1 DSL domain. Contains 6 EGF-like domains.
ドメイン	The DSL domain is required for binding to the Notch receptor.

画像




Flow Cytometry (Intracellular) - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) cells transfected with DLL3 expression construct containing a myc-His-tag[®] labelling DLL3 with **ab229902** at 1/600 dilution (Right) compared with a Rabbit monoclonal IgG (**ab172730**, Left) isotype control.

Goat anti rabbit IgG (Alexa Fluor[®]488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.

Cells were stained with rabbit IgG (Left) or **ab229902** (Right). Then stained with anti-myc-tag conjugated to Alexa Fluor[®]647.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab229902**).




Immunocytochemistry/ Immunofluorescence - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) cells labelling DLL3 with **ab229902** at 1/100 dilution, followed by **ab150080** AlexaFluor[®]594 Goat anti-Rabbit secondary antibody at 1/500 dilution (Green). An anti-myc-tag mAb (Alexa Fluor[®] 488 Conjugate) was used at 1/100 dilution. Confocal image showing positive staining in HEK-293T cells transfected with DLL3 expression construct containing a myc-His-tag[®] is observed. The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150080** AlexaFluor[®]594 Goat anti-Rabbit secondary at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab229902**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Immunohistochemical analysis of paraffin-embedded human small cell lung cancer tissue labeling DLL3 with **ab229902** at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on human small cell lung cancer (PMID: 28487384, 30397180) is observed.


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.

The section was incubated with **ab229902** for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab229902**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Immunohistochemical analysis of paraffin-embedded human glioma tissue labeling DLL3 with **ab229902** at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on human glioma (PMID: 30397180) is observed.


The section was incubated with **ab229902** for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab229902**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Immunohistochemical analysis of paraffin-embedded mouse embryonic brain of day 14.5 tissue labeling DLL3 with [ab229902](#) at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)). Positive staining on mouse embryonic brain of day 14.5 (PMID: 19389376) is observed.


The section was incubated with [ab229902](#) for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab229902](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Immunohistochemical analysis of paraffin-embedded human cerebrum tissue labeling DLL3 with [ab229902](#) at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)). **Negative control:** no staining on human cerebrum (PMID: 30397180) is observed.


The section was incubated with [ab229902](#) for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab229902](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Immunohistochemical analysis of paraffin-embedded mouse cerebrum tissue labeling DLL3 with **ab229902** at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). **Negative control:** no staining on mouse cerebrum (PMID: 30397180) is observed.


The section was incubated with **ab229902** for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab229902**).



Immunoprecipitation - Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

DLL3 was immunoprecipitated from 0.35 mg HEK-293T (human embryonic kidney epithelial cell) transfected with DLL3 expression construct containing a myc-His-tag® whole cell lysate with **ab229902** at 1/30 dilution. Western blot was performed on the immunoprecipitate using **ab229902** 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used as the secondary antibody at 1/5000 dilution.

Lane 1: HEK-293T (human embryonic kidney epithelial cell) transfected with DLL3 expression construct containing a myc-His-tag® whole cell lysate 10µg

Lane 2: **ab229902** IP in HEK-293T transfected with DLL3 expression construct containing a myc-His-tag® whole cell lysate (Input).

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab229902** in HEK-293T transfected with DLL3 expression construct containing a myc-His-tag® whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 15 seconds.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab229902**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-DLL3 antibody [EPR22592-18] - BSA and Azide free (ab255694)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors