

Anti-E tag antibody ab3397

★★★★★ [2 Abreviews](#) [11 References](#) [画像数 2](#)

製品の概要

製品名	Anti-E tag antibody
製品の詳細	Rabbit polyclonal to E tag
由来種	Rabbit
アプリケーション	適用あり: ICC/IF
種交差性	交差種: Species independent
免疫原	Synthetic peptide corresponding to E tag conjugated to keyhole limpet haemocyanin. Run BLAST with ExPASy Run BLAST with NCBI
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide Constituent: 99.91% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab3397の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
ICC/IF		Use at an assay dependent concentration.

ターゲット情報

関連性

E tag is a peptide epitope tag, sequence GAPVPYPDPLEPR.


画像



Immunofluorescent analysis of 4% paraformaldehyde (PFA)- fixed, permeabilized with 0.1% Triton X-100 in CHO cells transfected with 12-tags constructs (CHO-12 tags) labelling E tag with ab3397 at 1/1000 dilution, followed by Donkey Anti-rabbit IgG (H&L), FITC conjugated antibody at 1/100 dilution (Green) or Donkey Anti-rabbit IgG (H&L), Alexa Fluor594 antibody at 1/500 dilution (Red).

Nucleus was counterstained with DAPI (Blue).

Images also show parallel staining in parental CHO cell line.



ab3397 staining E tag in cultured human HUVEC cells by Immunocytochemistry. Cells were fixed with paraformaldehyde, permeabilized with 0.5% Triton x100 and blocking with 1% BSA was done for 45 minutes at 25°C. Samples were incubated with primary antibody (1/100: in 1% BSA/PBS) for 16 hours at 4°C. An FITC-conjugated donkey polyclonal to rabbit IgG was used at dilution at 1/200 as secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors