

Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody ab34711

8 References **画像数 4**

製品の概要

製品名	Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody
製品の詳細	Rabbit polyclonal to E1 Ubiquitin Activating Enzyme 1/UBA1
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rabbit, Dog 
免疫原	Fusion protein corresponding to Human E1 Ubiquitin Activating Enzyme 1/UBA1. Database link: P22314
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 0.42% Potassium phosphate, 0.87% Sodium chloride
精製度	IgG fraction
特記事項(精製)	The product was purified from monospecific antiserum by a multi step procedure.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab34711の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/5000. Detects a band of approximately 118 kDa (predicted molecular weight: 118 kDa).
IHC-P		Use a concentration of 2 - 20 µg/ml.
ICC/IF		Use a concentration of 5 µg/ml.

ターゲット情報

機能	Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding an ubiquitin-E1 thioester and free AMP.
パスウェイ	Protein modification; protein ubiquitination.
関連疾患	Defects in UBA1 are the cause of spinal muscular atrophy X-linked type 2 (SMA2) [MIM:301830]; also known as X-linked lethal infantile spinal muscular atrophy, distal X-linked arthrogryposis multiplex congenita or X-linked arthrogryposis type 1 (AMCX1). Spinal muscular atrophy refers to a group of neuromuscular disorders characterized by degeneration of the anterior horn cells of the spinal cord, leading to symmetrical muscle weakness and atrophy. SMA2 is a lethal infantile form presenting with hypotonia, areflexia, and multiple congenital contractures.
配列類似性	Belongs to the ubiquitin-activating E1 family.
翻訳後修飾	ISGylated.

画像



All lanes : Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody (ab34711) at 1 µg/ml

- Lane 1 :** Human brain tissue lysate - total protein ([ab29466](#))
- Lane 2 :** Human liver tissue lysate - total protein ([ab29889](#))
- Lane 3 :** HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate
- Lane 4 :** MCF7 (Human breast adenocarcinoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg/ml per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ([ab97080](#)) at 1/5000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 118 kDa

Observed band size: 118 kDa


Additional bands at: 35 kDa, 53 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 1 minute




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody (ab34711)

This image shows human lung tissue stained with ab34711 at 10µg/ml. In many cells apunctate nuclear staining was observed. Other cells showed both cytoplasmic and nuclear staining.



Immunocytochemistry/ Immunofluorescence - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody (ab34711)

ICC/IF image of ab34711 stained MCF7 cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab34711, 5µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody (ab34711)

ab34711 (4µg/ml) staining E1 Ubiquitin activating enzyme in human testis using an automated system (DAKO Autostainer Plus). Using this protocol there is nuclear and weak cytoplasmic staining. Sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffer EDTA pH 9.0 in a DAKO PT link. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako envision flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that, for manual staining, optimization of primary antibody concentration and incubation time is recommended. Signal amplification may be required.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors