abcam

Product datasheet

Anti-ENTPD5 antibody [EPR3783] ab92542

יעלאעבע RabMAb

1 References 画像数 2

製品の概要

製品名 Anti-ENTPD5 antibody [EPR3783]

製品の詳細 Rabbit monoclonal [EPR3783] to ENTPD5

由来種 Rabbit

アプリケーション **適用あり:** WB

適用なし: Flow Cyt,ICC/IF,IHC-P or IP

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

Synthetic peptide within Human ENTPD5 aa 100-200. The exact sequence is proprietary. 免疫原

ポジティブ・コントロール Human fetal liver, Hela and LNCaP lysates.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.

バッファー pH: 7.20

Preservative: 0.05% Sodium azide

Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue

culture supernatant

精製度 Protein A purified

ポリ/モノ モノクローナル クローン名 **EPR3783**

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab92542の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/5000. Predicted molecular weight: 47 kDa.


追加情報

Is unsuitable for Flow Cyt,ICC/IF,IHC-P or IP.

ターゲット情報

機能	Uridine diphosphatase (UDPase) that promotes protein N-glycosylation and ATP level regulation. UDP hydrolysis promotes protein N-glycosylation and folding in the endoplasmic reticulum, as well as elevated ATP consumption in the cytosol via an ATP hydrolysis cycle. Together with CMPK1 and AK1, constitutes an ATP hydrolysis cycle that converts ATP to AMP and results in a compensatory increase in aerobic glycolysis. Also hydrolyzes GDP and IDP but not any other nucleoside di-, mono- or triphosphates, nor thiamine pyrophosphate. Plays a key role in the AKT1-PTEN signaling pathway by promoting glycolysis in proliferating cells in response to phosphoinositide 3-kinase (PI3K) signaling.	
組織特異性	Expressed in adult liver, kidney, prostate, testis and colon. Much weaker expression in other tissues.	
パスウェイ	Protein modification; protein glycosylation.	
配列類似性	Belongs to the GDA1/CD39 NTPase family.	
翻訳後修飾	N-glycosylated; high-mannose type.	
細胞内局在	Endoplasmic reticulum lumen.	

画像



All lanes : Anti-ENTPD5 antibody [EPR3783] (ab92542) at 1/1000 dilution

Lane 1: Human fetal liver lysate

Lane 2 : Hela cell lysate


Lane 3 : LNCaP cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP conjugated Goat anti-Rabbit lgG at 1/2000 dilution

Predicted band size: 47 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors