

Anti-Factor H antibody [OX-24] ab118820

6 References [画像数 2](#)

製品の概要

製品名	Anti-Factor H antibody [OX-24]
製品の詳細	Mouse monoclonal [OX-24] to Factor H
由来種	Mouse
アプリケーション	適用あり: ELISA, WB
種交差性	交差種: Human
免疫原	Full length native protein (purified) corresponding to Human Factor H.
ポジティブ・コントロール	WB: Human plasma, Human serum, Purified Factor H protein
特記事項	<p>ab118820 has switched from ascites to TCS on 19th September 2019. Lot numbers higher than GR3258447 are from tissue culture supernatant.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium azide Constituent: PBS
精製度	Protein G purified
特記事項 (精製)	Purified from TCS.
ポリ/モノ	モノクローナル
クローン名	OX-24

アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab118820の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ELISA		Use a concentration of 10 µg/ml.
WB		Use at an assay dependent concentration. Predicted molecular weight: 139 kDa.

ターゲット情報

機能	Factor H functions as a cofactor in the inactivation of C3b by factor I and also increases the rate of dissociation of the C3bBb complex (C3 convertase) and the (C3b)NBB complex (C5 convertase) in the alternative complement pathway.
組織特異性	Expressed by the liver and secreted in plasma.
関連疾患	<p>Genetic variations in CFH are associated with basal laminar drusen (BLD) [MIM:126700]; also known as drusen of Bruch membrane or cuticular drusen or grouped early adult-onset drusen. Drusen are extracellular deposits that accumulate below the retinal pigment epithelium on Bruch membrane. Basal laminar drusen refers to an early adult-onset drusen phenotype that shows a pattern of uniform small, slightly raised yellow subretinal nodules randomly scattered in the macula. In later stages, these drusen often become more numerous, with clustered groups of drusen scattered throughout the retina. In time these small basal laminar drusen may expand and ultimately lead to a serous pigment epithelial detachment of the macula that may result in vision loss.</p> <p>Defects in CFH are the cause of complement factor H deficiency (CFH deficiency) [MIM:609814]. CFH deficiency determines uncontrolled activation of the alternative complement pathway with consumption of C3 and often other terminal complement components. It is associated with a number of renal diseases with variable clinical presentation and progression, including membranoproliferative glomerulonephritis and atypical hemolytic uremic syndrome. CFH deficiency patients may show increased susceptibility to meningococcal infections.</p> <p>Defects in CFH are a cause of susceptibility to hemolytic uremic syndrome atypical type 1 (AHUS1) [MIM:235400]. An atypical form of hemolytic uremic syndrome. It is a complex genetic disease characterized by microangiopathic hemolytic anemia, thrombocytopenia, renal failure and absence of episodes of enterocolitis and diarrhea. In contrast to typical hemolytic uremic syndrome, atypical forms have a poorer prognosis, with higher death rates and frequent progression to end-stage renal disease. Note=Susceptibility to the development of atypical hemolytic uremic syndrome can be conferred by mutations in various components of or regulatory factors in the complement cascade system. Other genes may play a role in modifying the phenotype.</p> <p>Genetic variation in CFH is associated with age-related macular degeneration type 4 (ARMD4) [MIM:610698]. ARMD is a multifactorial eye disease and the most common cause of irreversible vision loss in the developed world. In most patients, the disease is manifest as</p>

ophthalmoscopically visible yellowish accumulations of protein and lipid (known as drusen) that lie beneath the retinal pigment epithelium and within an elastin-containing structure known as Bruch membrane.


配列類似性

Contains 20 Sushi (CCP/SCR) domains.

細胞内局在

Secreted.

画像



All lanes : Anti-Factor H antibody [OX-24] (ab118820) at 1 µg/ml

Lane 1 : Human serum diluted 1/100 at 10 µl

Lane 2 : Human plasma diluted 1/100 at 10 µl

Lane 3 : Purified Factor H protein at 0.5 µg

Lane 4 : Purified Factor H protein at 0.1 µg


Performed under reducing conditions.

Predicted band size: 139 kDa

Observed band size: 170 kDa

This blot was produced using a 4-12% Bis-tris under the MOPS buffer system. The gel was run at 200V for 55 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% milk before being incubated with ab118820 overnight at 4°C at a 1µg/ml concentration. Antibody binding was detected using Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed ([ab216772](#)) at 1/20000 dilution for 1 hour at room temperature before imaging.

This image was generated using the ascites version of the product.



96-well microtitre plates were coated overnight at 4°C with recombinant human C3, C3b, iC3b, and Factor H proteins, in duplicate at a concentration of 1µg/mL. Plates were blocked with 1% BSA in PBS-T (0.1% Tween®) for 1 hour before incubation with a 10-step 4x serial dilution of ab118820 from 10µg/mL for 1 hour at room temperature. Antibody binding was detected with Goat Anti-Mouse IgG H&L (HRP) (**ab6789**) secondary antibody at a 1 in 10000 dilution for 1 hour at room temperature. Plates were incubated with TMB ELISA substrate for 7 minutes prior to being stopped with Stop solution and absorbance measured at 450nm.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors