# abcam

## Product datasheet

## Anti-FAK antibody [EP1831Y] - BSA and Azide free ab247481



יעלטעבע RabMAb

## 画像数 2

#### 製品の概要

製品名 Anti-FAK antibody [EP1831Y] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EP1831Y] to FAK - BSA and Azide free

由来種 Rabbit

アプリケーション 適用あり: IHC-P, WB

適用なし: №

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

特記事項 ab247481 is the carrier-free version of ab76496.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

**バッファー** pH: 7.2

Constituent: PBS

キャリア・フリー はい

精製度 Protein A purified

**ポリ/モノ** モノクローナル **クローン名** EP1831Y

アイソタイプ IgG

## アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab247481の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Predicted molecular weight: 119 kDa.

追加情報 Is unsuitable for IP.

## ターゲット情報

機能 Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility,

proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Microtubule-induced dephosphorylation at Tyr-397 is crucial for the induction of focal adhesion disassembly. Plays a potential role in oncogenic transformations resulting in increased

kinase activity.

組織特異性 Expressed in all organs tested, in lymphoid cell lines, but most abundantly in brain.

**配列類似性** Belongs to the protein kinase superfamily. Tyr protein kinase family. FAK subfamily.

Contains 1 FERM domain.

Contains 1 protein kinase domain.

ドメイン The first Pro-rich domain interacts with the SH3 domain of CRK-associated substrate (BCAR1)

and CASL.


The carboxy-terminal region is the site of focal adhesion targeting (FAT) sequence which

mediates the localization of FAK1 to focal adhesions.

翻訳後修飾 Phosphorylated on 6 tyrosine residues upon activation. Microtubule-induced dephosphorylation at

Tyr-397 could be catalyzed by PTPN11 and regulated by ZFYVE21. Dephosphorylated by

#### 画像



Western blot - Anti-FAK antibody [EP1831Y] - BSA and Azide free (ab247481)

**All lanes :** Anti-FAK antibody [EP1831Y] (ab76496) at 1/500 dilution

Lane 1 : HeLa cell lysate

Lane 2 : A431 cell lysate

Lane 3: Wild-type HEK-293T cell lysate

Lane 4: PTK2 knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 119 kDa Observed band size: 119 kDa

This data was developed using the same antibody clone in a different buffer formulation (ab76496).

**Lanes 1 - 4:** Merged signal (red and green). Green - <u>ab76496</u> observed at 119 kDa. Red - loading control, <u>ab8245</u> observed at 37 kDa.

ab76496 was shown to react with FAK in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line ab255421 (knockout cell lysate ab263766) was used. Wild-type and FAK knockout samples were subjected to SDS-PAGE. ab76496 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) were incubated overnight at 4°C at 1 in 500 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors