

Anti-FGD4 antibody ab97785

4 References [画像数 3](#)

医薬用外劇物

製品の概要

製品名	Anti-FGD4 antibody
製品の詳細	Rabbit polyclonal to FGD4
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF
種交差性	交差種: Human
免疫原	Recombinant fragment, corresponding to a sequence within amino acids 49-297 of Human FGD4 (NP_640334).
ポジティブ・コントロール	293T, A431, H1299, HeLa, HepG2, MOLT4 and Raji whole cell lysates; HeLa cells; SG xenograft tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	<p>pH: 7.00</p> <p>Preservative: 0.01% Thimerosal (merthiolate)</p> <p>Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)</p>
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab97785の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/3000. Predicted molecular weight: 87 kDa.
IHC-P		Use at an assay dependent concentration.
ICC/IF		1/100 - 1/200.

ターゲット情報

機能	Activates CDC42, a member of the Ras-like family of Rho-and Rac proteins, by exchanging bound GDP for free GTP. Plays a role in regulating the actin cytoskeleton and cell shape. Activates MAPK8.
組織特異性	Expressed in different tissues, including brain, cerebellum, peripheral nerve, skeletal muscle, heart, uterus, placenta and testis.
関連疾患	Charcot-Marie-Tooth disease 4H (CMT4H) [MIM:609311]: A recessive demyelinating form of Charcot-Marie-Tooth disease, a disorder of the peripheral nervous system, characterized by progressive weakness and atrophy, initially of the peroneal muscles and later of the distal muscles of the arms. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathies (designated CMT1 when they are dominantly inherited) and primary peripheral axonal neuropathies (CMT2). Demyelinating neuropathies are characterized by severely reduced nerve conduction velocities (less than 38 m/sec), segmental demyelination and remyelination with onion bulb formations on nerve biopsy, slowly progressive distal muscle atrophy and weakness, absent deep tendon reflexes, and hollow feet. By convention autosomal recessive forms of demyelinating Charcot-Marie-Tooth disease are designated CMT4. Note=The disease is caused by mutations affecting the gene represented in this entry.
配列類似性	Contains 1 DH (DBL-homology) domain. Contains 1 FYVE-type zinc finger. Contains 2 PH domains.
ドメイン	The part of the protein spanning the actin filament-binding domain together with the DH domain and the first PH domain is necessary and sufficient for microspike formation. Activation of MAPK8 requires the presence of all domains with the exception of the actin filament-binding domain.
細胞内局在	Cytoplasm > cytoskeleton. Cell projection > filopodium. Concentrated in filopodia and poorly detected at lamellipodia. Binds along the sides of actin fibers.

画像





Western blot - Anti-FGD4 antibody (ab97785)

Anti-FGD4 antibody (ab97785) at 1/1000 dilution + MOLT4 whole cell lysate at 30 µg


Predicted band size: 87 kDa

7.5% SDS-PAGE.



Immunocytochemistry/ Immunofluorescence - Anti-FGD4 antibody (ab97785)

ab97785 at 1/200 dilution staining FGD4 in HeLa cells by Immunofluorescence, Paraformaldehyde fixed. Lower image is merged with DNA probe.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FGD4 antibody (ab97785)

ab97785 at 1/100 dilution staining FGD4 in SG xenograft by Immunohistochemistry, Paraffin-embedded tissue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors