abcam

Product datasheet

FITC Anti-Carcino Embryonic Antigen CEA antibody [CB30] ab106739

2 References

製品の概要

製品名 FITC Anti-Carcino Embryonic Antigen CEA antibody [CB30]

製品の詳細 FITC Mouse monoclonal [CB30] to Carcino Embryonic Antigen CEA

由来種 Mouse

標識 FITC. Ex: 493nm, Em: 528nm

アプリケーション 適用あり: Flow Cyt

種交差性 交差種: Human, Monkey

免疫原 Full length protein. Human hepatoma cell line

ポジティブ・コントロール Strongly stains tumor cells present in cancerous colon tissue. Only very weak staining of normal

tissue.

特記事項
The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.40

Preservative: 0.1% Sodium azide Constituents: PBS, 0.5% BSA

精製度 Size exclusion

特記事項(精製) Purified fraction from size exclusion chromatography.

ポリ/モノ モノクローナル

クローン名 CB30

1

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab106739の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration. (leucocytes) in 100 ul. <u>ab91356</u> - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Cell surface glycoprotein that plays a role in cell adhesion and in intracellular signaling. Receptor for E.coli Dr adhesins.	
組織特異性	Found in adenocarcinomas of endodermally derived digestive system epithelium and fetal colon.	
配列類似性	Belongs to the immunoglobulin superfamily. CEA family. Contains 7 lg-like (immunoglobulin-like) domains.	
翻訳後修飾	Complex immunoreactive glycoprotein with a MW of 180 kDa comprising 60% carbohydrate.	
細胞内局在	Cell membrane.	

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors