abcam

Product datasheet

FITC Anti-CD79b antibody [AT107-2] ab33296

1 References 画像数 1

製品の概要

製品名 FITC Anti-CD79b antibody [AT107-2]

製品の詳細 FITC Rat monoclonal [AT107-2] to CD79b

由来種 Rat

標識 FITC. Ex: 493nm, Em: 528nm

特異性 Recognises an intracellular region of human CD79b We have data to indicate that this antibody

may not cross react with Chicken. However, this has not been conclusively tested and expression

levels may vary in certain cell lines/tissues.

アプリケーション 適用あり: Flow Cyt (Intra)

種交差性 交差種: Rat

免疫原 Synthetic peptide corresponding to an intracellular region of human CD79b.

ポジティブ・コントロール Flow Cyt (Intra): Rat peripheral blood lymphocytes.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.40

Preservative: 0.09% Sodium azide

Constituents: PBS, 1% BSA

精製度 Protein G purified

ポリ/モノ モノクローナル

クローン名 AT107-2

₹II-7 NS1

1

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab33296の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		

追加情報 Flow Cyt: Use 10µl for 10⁶ cells in 100µl.

Note: Membrane permeabilisation is required for this application.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

機能 Required in cooperation with CD79A for initiation of the signal transduction cascade activated by

the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Enhances phosphorylation of CD79A, possibly by recruiting kinases which phosphorylate CD79A or by recruiting proteins which bind to CD79A and

protect it from dephosphorylation.

組織特異性 B-cells.

関連疾患 Defects in CD79B are the cause of agammaglobulinemia type 6 (AGM6) [MIM:612692]. It is a

primary immunodeficiency characterized by profoundly low or absent serum antibodies and low or

absent circulating B cells due to an early block of B-cell development. Affected individuals

develop severe infections in the first years of life.

配列類似性 Contains 1 lg-like V-type (immunoglobulin-like) domain.

Contains 1 ITAM domain.


翻訳後修飾 Phosphorylated on tyrosine upon B-cell activation.

細胞内局在 Cell membrane. Following antigen binding, the BCR has been shown to translocate from

detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through

the complex can also occur outside lipid rafts.

画像



Staining of rat peripheral blood lymphocytes with ab33296 after permeabilisation.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors