abcam

Product datasheet

FITC Anti-IL-2 Receptor alpha antibody [MEM-181] ab18275

製品の概要

製品名 FITC Anti-IL-2 Receptor alpha antibody [MEM-181]

製品の詳細 FITC Mouse monoclonal [MEM-181] to IL-2 Receptor alpha

由来種 Mouse

標識 FITC. Ex: 493nm, Em: 528nm

アプリケーション 適用あり: Flow Cyt **種交差性** 交差種: Human

免疫原 Tissue, cells or virus corresponding to Human IL-2 Receptor alpha.

ポジティブ・コントロール Human blood cells.

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.40

Preservative: 0.097% Sodium azide

Constituents: PBS, BSA

精製度 Protein G purified

ポリ/モノ モノクローナル **クローン名** MEM-181

アイソタイプ IgG1

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab18275の使用に適用されます

dependent type 10 (IDDM10) [MIM:601942]. A multifactorial disorder of glucose homeostasis that

osmotic diuresis and secondary thirst. These derangements result in long-term complications that

is characterized by susceptibility to ketoacidosis in the absence of insulin therapy. Clinical fetaures are polydipsia, polyphagia and polyuria which result from hyperglycemia-induced

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項	
Flow Cyt			
追加情報	Flow Cyt: Use at a concentration of 20 μ l/100 μ l of whole human blood.		
	Not tested in other applications.		
	Optimal dilutions/concer	Optimal dilutions/concentrations should be determined by the end user.	
ターゲット情報			
機能	Receptor for interleukin-	Receptor for interleukin-2.	
関連疾患	Genetic variations in IL2RA are associated with susceptibility to diabetes mellitus insulin-		

Contains 2 Sushi (CCP/SCR) domains.

affect the eyes, kidneys, nerves, and blood vessels.

細胞内局在 Membrane.

配列類似性

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors