abcam

Product datasheet

FITC Anti-Listeria monocytogenes antibody ab68592

1 References

製品の概要

製品名 FITC Anti-Listeria monocytogenes antibody

製品の詳細 FITC Rabbit polyclonal to Listeria monocytogenes

由来種 Rabbit

標識 FITC. Ex: 493nm, Em: 528nm

特異性 This antibody reacts with Listeria sp. in bacterial and infected tissue samples.

アプリケーション 適用あり: IHC-Fr, ICC/IF

種交差性 交差種: Listeria monocytogenes

免疫原 Tissue, cells or virus corresponding to Listeria monocytogenes. Listeria monocytogenes; ATCC

strain 43251.

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.2

Preservative: 0.1% Sodium azide Constituents: 1% BSA, PBS

精製度 Affinity purified

ポリ/モノ ポリクローナル

アイソタイプ lqG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab68592の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		1/10.
ICC/IF		1/10.

ターゲット情報

関連性

The genus Listeria comprises six species: L. monocytogenes, L. innocua, L. welshimeri, L. seeligeri, L. ivanovii and L. grayi. Listeria monocytogenes, the most commonly isolated pathogenic member, is associated with a wide spectrum of human and animal diseases. In the smear from the original tissue, L. monocytogenes may appear as gram-positive coccobacilli that may be confused with Streptococcus agalactiae (group B), enterococci, or Corynebacterium spp. Listeria is differentiated from streptococci by a positive catalase test. L. monocytogenes is the only species of the genus Listeria that has been clearly documented as a pathogen for humans. The forms of disease caused by this organism are myriad and age-related. The most common clinical manifestations are meningitis and septicemia. Listeria monocytogenes, a food-borne intracellular animal and human pathogen, interacts with infected host cells both prior to entry and during the intracellular phase of infection.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors