

Anti-FKBP51 antibody [EPR6617] ab126715

KO 評価済 リコンビナント RabMAb


★★★★★ 1 Abreviews 9 References 画像数 11

製品の概要

製品名	Anti-FKBP51 antibody [EPR6617]
製品の詳細	Rabbit monoclonal [EPR6617] to FKBP51
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF
種交差性	交差種: Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Human colon, Rat stomach and Human prostatic hyperplasia tissue. WB: Human testis tissue; Jurkat, HepG2, Caco-2 and HeLa whole cell lysate (ab150035). Wild-type HAP1 whole cell lysate. ICC/IF: Jurkat cells. Flow Cyt (intra): HeLa cells. IP: Jurkat cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
解離定数 (K _D 値)	K _D = 7.10 x 10 ⁻¹¹ M



バッファー	pH: 7.20 Preservative: 0.01% Sodium azide
-------	--

	Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR6617
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab126715の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/10 - 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (1)	1/1000 - 1/10000. Predicted molecular weight: 51 kDa.
IP		1/10 - 1/100.
IHC-P		1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/50 - 1/100.
ICC/IF		1/100 - 1/250.

ターゲット情報

機能	Interacts with functionally mature heterooligomeric progesterone receptor complexes along with HSP90 and TEBP.
組織特異性	Widely expressed, enriched in testis compared to other tissues.
配列類似性	Contains 2 PPlase FKBP-type domains. Contains 3 TPR repeats.
翻訳後修飾	Phosphorylated upon DNA damage, probably by ATM or ATR.
細胞内局在	Cytoplasm. Nucleus.

画像



Western blot - Anti-FKBP51 antibody [EPR6617]
(ab126715)

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: FKBP51 knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)

Lane 4: HepG2 whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab126715 observed at 51 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab126715 was shown to recognize FKBP51 in wild-type cells as signal was lost at the expected MW in FKBP51 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and FKBP51 knockout samples were subjected to SDS-PAGE. Ab126715 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-FKBP51 antibody [EPR6617]
(ab126715)

All lanes : Anti-FKBP51 antibody [EPR6617] (ab126715) at 1/10000 dilution (Purified)

Lane 1 : Jurkat (Human T cell leukemia T lymphocyte) whole cell lysates at 15 µg


Lane 2 : Rat spleen lysates at 15 µg

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 51 kDa

Observed band size: 51 kDa



Immunoprecipitation - Anti-FKBP51 antibody
[EPR6617] (ab126715)

ab126715 (purified) at 1:20 dilution (2µg) immunoprecipitating FKBP51 in Jurkat whole cell lysate.


Lane 1 (input): Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10µg

Lane 2 (+): ab126715 & Jurkat whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab126715 in Jurkat whole cell lysate


For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FKBP51 antibody
[EPR6617] (ab126715)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat stomach tissue sections labeling FKBP51 with purified ab126715 at 1:250 dilution (1.156 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FKBP51 antibody [EPR6617] (ab126715)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human prostatic hyperplasia tissue sections labeling FKBP51 with purified ab126715 at 1:250 dilution (1.156 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunocytochemistry/ Immunofluorescence - Anti-FKBP51 antibody [EPR6617] (ab126715)


Immunocytochemistry/ Immunofluorescence analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling FKBP51 with purified ab126715 at 1:100 dilution (2.9 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

Alexa Fluor® 488 (**ab198978**) and Alexa Fluor® 647 (**ab198979**) conjugated versions are available for this clone.



Flow Cytometry (Intracellular) - Anti-FKBP51 antibody [EPR6617] (ab126715)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling FKBP51 with purified ab126715 at 1/20 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue). Alexa Fluor® 488 (**ab198978**) and Alexa Fluor® 647 (**ab198979**) conjugated versions are available for this clone.



Western blot - Anti-FKBP51 antibody [EPR6617] (ab126715)

All lanes : Anti-FKBP51 antibody [EPR6617] (ab126715) at 1/1000 dilution


Lane 1 : Human testis cell lysate

Lane 2 : Jurkat cell lysate

Lane 3 : Caco-2 cell lysate

Lane 4 : HeLa cell lysate


Predicted band size: 51 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FKBP51 antibody [EPR6617] (ab126715)

ab126715, at 1/50 dilution, stains FKBP51 in paraffin embedded human colon tissue by immunohistochemistry.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



OL-RD Scanning - Anti-FKBP51 antibody [EPR6617]
(ab126715)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a
recombinant antibody?



Research with
confidence
Consistent and
reproducible results



Long-term and
scalable supply
Recombinant
technology



Success from the
first experiment
Confirmed
specificity



Ethical standards
compliant
Animal-free
production

Anti-FKBP51 antibody [EPR6617] (ab126715)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors