

### Anti-FTO antibody [5-2H10] ab92821

KO 評価済

★★★★☆ **3 Abreviews** **42 References** **画像数 4**

#### 製品の概要

製品名	Anti-FTO antibody [5-2H10]
製品の詳細	Mouse monoclonal [5-2H10] to FTO
由来種	Mouse
アプリケーション	<b>適用あり:</b> WB, ICC/IF, Flow Cyt
種交差性	<b>交差種:</b> Rat, Human <b>交差が予測される動物種:</b> Sheep, Horse, Cow, Dog, Non human primates 
免疫原	Synthetic peptide corresponding to Human FTO (N terminal) conjugated to keyhole limpet haemocyanin. Database link: <a href="#">Q9C0B1</a>
ポジティブ・コントロール	WB: Rat testes nuclear lysate. ICC/IF: Neuronal progenitor cells. Flow Cyt: SH-SY5Y cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	Constituents: 0.238% HEPES, 0.01% BSA, 50% Glycerol, 0.87% Sodium chloride
精製度	Protein G purified
特記事項(精製)	Protein G purified culture supernatant.
ポリ/モノ	モノクローナル
クローン名	5-2H10
アイソタイプ	IgG2a

## アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab92821の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 58 kDa (predicted molecular weight: 58 kDa).
ICC/IF	★★★★★ (1)	1/100.
Flow Cyt		Use 1µg for 10 <sup>6</sup> cells. <b>ab170191</b> - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

## ターゲット情報

**機能** Dioxygenase that repairs alkylated DNA and RNA by oxidative demethylation. Has highest activity towards single-stranded RNA containing 3-methyluracil, followed by single-stranded DNA containing 3-methylthymine. Has low demethylase activity towards single-stranded DNA containing 1-methyladenine or 3-methylcytosine. Has no activity towards 1-methylguanine. Has no detectable activity towards double-stranded DNA. Requires molecular oxygen, alpha-ketoglutarate and iron. Contributes to the regulation of the global metabolic rate, energy expenditure and energy homeostasis. Contributes to the regulation of body size and body fat accumulation.

**組織特異性** Ubiquitously expressed, with relatively high expression in adrenal glands and brain; especially in hypothalamus and pituitary.


**関連疾患** Defects in FTO are the cause of growth retardation developmental delay coarse facies and early death (GRDDCFED) [MIM:612938]. The disease consists of a severe children multiple congenital anomaly syndrome with death by the age of 3 years. All affected individuals had postnatal growth retardation, microcephaly, severe psychomotor delay, functional brain deficits, and characteristic facial dysmorphism. In some patients, structural brain malformations, cardiac defects, genital anomalies, and cleft palate were also observed.

**配列類似性** Belongs to the fto family.

**ドメイン** The 3D-structure of the Fe2OG dioxygenase domain is similar to that of the Fe2OG dioxygenase domain found in the bacterial DNA repair dioxygenase alkB and its mammalian orthologs, but sequence similarity is very low. As a consequence, the domain is not detected by protein signature databases.

**細胞内局在** Nucleus.

## 画像



Western blot - Anti-FTO antibody [5-2H10]  
(ab92821)

**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)


**Lane 2:** FTO knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** HEK293 whole cell lysate (20 µg)

**Lane 4:** MOLT4 whole cell lysate (20 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab92821 observed at 58 kDa. Red - loading control, **ab181602**, observed at 37 kDa.

ab92821 was shown to specifically recognize FTO in wild-type HAP1 cells along with additional cross reactive bands. No band was observed when FTO knockout samples were examined. Wild-type and FTO knockout samples were subjected to SDS-PAGE. ab92821 and **ab181602** (Rabbit anti-GAPDH loading control) were incubated overnight at 4°C at a 1/1000 dilution and 1/10,000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed (**ab216772**) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed (**ab216777**) secondary antibodies at 1/10,000 dilution for 1 hour at room temperature before imaging.




Flow Cytometry - Anti-FTO antibody [5-2H10]  
(ab92821)

Overlay histogram showing SH-SY5Y cells stained with ab92821 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab92821, 1 µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIG2A] (**ab91361**, 1 µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in SH-SY5Y cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Anti-FTO antibody [5-2H10] (ab92821)

ICC/IF image of ab92821 stained PC12 cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab92821, 5µg/ml) overnight at +4°C. The secondary antibody (green) was **ab96879**, DyLight® 488 goat anti-mouse IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Western blot - Anti-FTO antibody [5-2H10] (ab92821)

Anti-FTO antibody [5-2H10] (ab92821) at 1/1000 dilution + Rat testes lysate

**Predicted band size:** 58 kDa

**Observed band size:** 58 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors