

Anti-FUBP1/FBP antibody [EPR12327] ab181111

リコンビナント **RabMAb**

★★★★★ **2 Abreviews** **12 References** 画像数 **14**

製品の概要

製品名	Anti-FUBP1/FBP antibody [EPR12327]
製品の詳細	Rabbit monoclonal [EPR12327] to FUBP1/FBP
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IHC-P, ICC/IF
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Jurkat, HeLa, Raji and HepG2 whole cell lysate (ab7900), rat brain, mouse brain and mouse spleen tissue lysates. IHC-P: Human breast carcinoma and pancreas tissues. ICC/IF: HeLa and HepG2 cells. Flow Cyt (intra): Jurkat cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR12327
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab181111の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/60 - 1/90. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		1/2000. Detects a band of approximately 74 kDa (predicted molecular weight: 68 kDa). For unpurified use at 1/1000 - 1/20000.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .
ICC/IF	★★★★★ (1)	1/250 - 1/500.

ターゲット情報

機能	Regulates MYC expression by binding to a single-stranded far-upstream element (FUSE) upstream of the MYC promoter. May act both as activator and repressor of transcription.
配列類似性	Contains 4 KH domains.
翻訳後修飾	Ubiquitinated. This targets the protein for proteasome-mediated degradation.
細胞内局在	Nucleus.

画像



All lanes : Anti-FUBP1/GBP antibody [EPR12327] (ab181111) at 1/2000 dilution (purified)

Lane 1 : Mouse brain tissue lysate

Lane 2 : Mouse spleen tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

Predicted band size: 68 kDa

Observed band size: 74 kDa

Blocking and dilution buffer: 5% NFDM/TBST.



Western blot - Anti-FUBP1/FBP antibody
[EPR12327] (ab181111)

Anti-FUBP1/FBP antibody [EPR12327] (ab181111) at 1/5000
dilution (purified) + Rat brain tissue lysate at 20 µg


Secondary

Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000
dilution

Predicted band size: 68 kDa

Observed band size: 74 kDa

Blocking and dilution buffer: 5% NFDM/TBST.



Western blot - Anti-FUBP1/FBP antibody
[EPR12327] (ab181111)

Anti-FUBP1/FBP antibody [EPR12327] (ab181111) at 1/5000
dilution (purified) + HepG2 cell lysate at 20 µg


Secondary

Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000
dilution

Predicted band size: 68 kDa

Observed band size: 74 kDa

Blocking and dilution buffer: 5% NFDM/TBST.



All lanes : Anti-FUBP1/FBP antibody [EPR12327] (ab181111) at 1/20000 dilution (purified)

Lane 1 : Jurkat cell lysate

Lane 2 : HeLa cell lysate

Lane 3 : Raji cell lysate

Lane 4 : K562 cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

Predicted band size: 68 kDa


Observed band size: 74 kDa

Blocking and dilution buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FUBP1/GBP antibody [EPR12327] (ab181111)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labelling FUBP1/GBP with purified ab181111 at 1/500. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.




Immunocytochemistry/ Immunofluorescence - Anti-FUBP1/GBP antibody [EPR12327] (ab181111)

Immunocytochemistry/Immunofluorescence analysis of HeLa cells labelling FUBP1/GBP with purified ab181111 at 1/500. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.


Control 1: primary antibody (1/500) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).



Flow Cytometry (Intracellular) - Anti-FUBP1/GBP antibody [EPR12327] (ab181111)

Intracellular Flow Cytometry analysis of Jurkat cells labelling FUBP1 / GBP with purified ab181111 at 1/60 (red). Cells were fixed with 80% methanol. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



Western blot - Anti-FUBP1/FBP antibody
[EPR12327] (ab181111)


Anti-FUBP1/FBP antibody [EPR12327] (ab181111) at 1/5000
dilution (unpurified) + Jurkat cell lysate at 20 µg

Secondary

Goat anti-rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution

Predicted band size: 68 kDa

Observed band size: 74 kDa



Western blot - Anti-FUBP1/FBP antibody
[EPR12327] (ab181111)

All lanes : Anti-FUBP1/FBP antibody [EPR12327] (ab181111) at
1/20000 dilution (unpurified)

Lane 1 : HeLa cell lysate

Lane 2 : HepG2 cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-rabbit IgG, (H+L), Peroxidase conjugate at
1/1000 dilution

Predicted band size: 68 kDa


Observed band size: 74 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FUBP1/GBP antibody [EPR12327] (ab181111)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labeling FUBP1/GBP with unpurified ab181111 at a dilution of 1/500 followed by prediluted HRP Polymer for rabbit IgG. Counterstained with Hematoxylin.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FUBP1/GBP antibody [EPR12327] (ab181111)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human pancreas tissue labeling FUBP1/GBP with unpurified ab181111 at a dilution of 1/500 followed by prediluted HRP Polymer for rabbit IgG. Counterstained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-FUBP1/GBP antibody [EPR12327] (ab181111)

Immunocytochemistry/Immunofluorescent analysis of 4% paraformaldehyde-fixed HepG2 cells labeling FUBP1/GBP with unpurified ab181111 at 1/500 dilution (red) followed by Alexa Fluor® 555-conjugated goat anti rabbit IgG secondary antibody at 1/200 dilution. Counter stained with Dapi (blue).



Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed Jurkat cells labeling FUBP1/GBP with unpurified ab181111 at a dilution of 1/90 (pink), compared to a rabbit monoclonal IgG isotype control (green), followed by goat anti rabbit IgG (FITC) secondary antibody at a dilution of 1/150.

Why choose a recombinant antibody?

<p>Research with confidence Consistent and reproducible results</p>	<p>Long-term and scalable supply Recombinant technology</p>
<p>Success from the first experiment Confirmed specificity</p>	<p>Ethical standards compliant Animal-free production</p>

Anti-FUBP1/GBP antibody [EPR12327] (ab181111)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors