# abcam

### Product datasheet

## Anti-Galectin 7 antibody [EPR4287] - BSA and Azide free ab247704

יובעדער RabMAb

画像数6

#### 製品の概要

製品名 Anti-Galectin 7 antibody [EPR4287] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EPR4287] to Galectin 7 - BSA and Azide free

由来種 Rabbit

アプリケーション 適用あり: Flow Cyt (Intra), IHC-P, IP, ICC/IF, WB

種交差性 交差種: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール IP: HaCaT whole cell lysate. ICC/IF: HaCaT cells; Flow Cyt (intra): HaCaT; WB: Human skin

lysate, IHC: Human tonsil tissue.

特記事項 ab247704 is the carrier-free version of ab108623.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

1

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

**バッファー** pH: 7.2

Constituent: PBS

キャリア・フリー はい

精製度 Protein A purified

**ポリ/モノ** モノクローナル **クローン名** EPR4287

アイソタイプ lgG

#### アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab247704の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
IP		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 15 kDa (predicted molecular weight: 15 kDa).

## ターゲット情報

機能 Could be involved in cell-cell and/or cell-matrix interactions necessary for normal growth control.


Pro-apoptotic protein that functions intracellularly upstream of JNK activation and cytochrome c

release.

組織特異性 Mainly expressed in stratified squamous epithelium.

**配列類似性** Contains 1 galectin domain.

**細胞内局在** Cytoplasm. Nucleus. Secreted. May be secreted by a non-classical secretory pathway.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Galectin 7 antibody

[EPR4287] - BSA and Azide free (ab247704)

This data was developed using <u>ab108623</u>, the same antibody clone in a different buffer formulation.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human tonsil tissue sections labeling Galectin 7 with purified <u>ab108623</u> at 1/1500 dilution (0.35 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Rabbit specific IHC polymer detection kit HRP/DAB (<u>ab209101</u>) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-Galectin 7 antibody [EPR4287] - BSA and Azide free (ab247704)


Anti-Galectin 7 antibody [EPR4287] (ab108623) at 1/1000 dilution (Purified) + Human skin lysate

#### Secondary

Goat Anti-Rabbit lgG (HRP) with minimal cross-reactivity with human lgG at 1/2000 dilution


Predicted band size: 15 kDa

This data was developed using <u>ab108623</u>, the same antibody clone in a different buffer formulation.



Flow Cytometry (Intracellular) - Anti-Galectin 7 antibody [EPR4287] - BSA and Azide free (ab247704)


This data was developed using <u>ab108623</u>, the same antibody clone in a different buffer formulation. Intracellular Flow Cytometry analysis of HaCaT (Human skin keratinocyte) cells labeling Galectin 7 with Purified <u>ab108623</u> at 1/50 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit lgG (Alexa Fluor<sup>®</sup> 488 ,<u>ab150077</u>) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal lgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunocytochemistry/ Immunofluorescence - Anti-Galectin 7 antibody [EPR4287] - BSA and Azide free (ab247704)

This data was developed using <u>ab108623</u>, the same antibody clone in a different buffer formulation.

Immunocytochemistry analysis of HaCaT (Human skin keratinocyte) cells labeling Galectin 7 with Purified  $\underline{ab108623}$  at 1:1000 dilution (0.5 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit lgG (Alexa Fluor® 488, $\underline{ab150077}$ ) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunoprecipitation - Anti-Galectin 7 antibody [EPR4287] - BSA and Azide free (ab247704)

This data was developed using <u>ab108623</u>, the same antibody clone in a different buffer formulation.

Purified <u>ab108623</u> at 1/50 dilution ( $2\mu g$ ) immunoprecipitating Galectin 7 in HaCaT whole cell lysate.

Lane 1 (input): HaCaT (Human skin keratinocyte) whole cell lysate 10ug

Lane 2 (+): ab108623 + HaCaT whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of


ab108623 in HaCaT whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) (ab131366) (1/1000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: 15 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors