abcam

Product datasheet

Anti-Gli1 antibody [EPR4523] ab134906

ועלשעבע RabMAb

★★★★★ 2 Abreviews 30 References 画像数5

製品の概要

製品名 Anti-Gli1 antibody [EPR4523]

製品の詳細 Rabbit monoclonal [EPR4523] to Gli1

由来種 Rabbit

アプリケーション **適用あり:** WB

適用なし: Flow Cyt,ICC/IF,IHC-P or IP

種交差性 交差種: Human

免疫原 Synthetic peptide within Human Gli1 aa 1-100. The exact sequence is proprietary.

ポジティブ・コントロール RMS13 and A673 cell lysates; Human brain and testis tissues; A673 cells.

特記事項 This antibody was developed as part of a collaboration with the lab of Martin Fernandez-Zapico at

the Mayo Clinic.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at -20°C.

バッファー pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 40% Glycerol, 0.05% BSA, 59% PBS

精製度 Protein A purified

ポリ/モノ モノクローナル **ウローン名** EPR4523

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab134906の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★ ☆☆ <u>(2)</u>	1/100 - 1/1000. Predicted molecular weight: 118 kDa.

追加情報 Is unsuitable for Flow Cyt,ICC/IF,IHC-P or IP.

ターゲット情報

機能 Acts as a transcriptional activator. May regulate the transcription of specific genes during normal

development. May play a role in craniofacial development and digital development, as well as development of the central nervous system and gastrointestinal tract. Mediates SHH signaling and

thus cell proliferation and differentiation.

組織特異性 Testis, myometrium and fallopian tube. Also expressed in the brain with highest expression in the

cerebellum, optic nerve and olfactory tract.

配列類似性 Belongs to the GLI C2H2-type zinc-finger protein family.

Contains 5 C2H2-type zinc fingers.


翻訳後修飾 Phosphorylated in vitro by ULK3.

細胞内局在 Cytoplasm. Nucleus. Tethered in the cytoplasm by binding to SUFU. Activation and translocation

to the nucleus is promoted by interaction with STK36. Phosphorylation by ULK3 may promote

nuclear localization. Translocation to the nucleus is promoted by interaction with ZIC1.

画像



Western blot - Anti-Gli1 antibody [EPR4523] (ab134906)

All lanes : Anti-Gli1 antibody [EPR4523] (ab134906) at 1/1000 dilution

Lane 1: RMS 13 (human rhabdomyosarcoma fibroblast)
transfected with scrambled siRNA control whole cell lysate
Lane 2: RMS 13 (human rhabdomyosarcoma fibroblast)
transfected with siRNA specifically targeting Gli1 whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution

Predicted band size: 118 kDa Observed band size: 150 kDa

Exposure time: 180 seconds


Diluting and blocking buffer and concentration: 5% NFDM/TBST This blot was developed using a high sensitivity ECL substrate.

All lanes : Anti-Gli1 antibody [EPR4523] (ab134906) at 1/1000 dilution

Lane 1 : A673 (human muscle Ewing's Sarcoma) whole cell lysate Lane 2 : HepG2 (human hepatocellular carcinoma epithelial cell), whole cell lysate

Lane 3: RD (Human muscle rhabdomyosarcoma) whole cell lysate

Lysates/proteins at 20 µg per lane.



Western blot - Anti-Gli1 antibody [EPR4523] (ab134906)

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution

Predicted band size: 118 kDa Observed band size: 150 kDa

Exposure time: 180 seconds

Diluting and blocking buffer and concentration: 5% NFDM/TBST

This blot was developed using a high sensitivity ECL substrate.

The expression level of Gli1 in HepG2 and RD are consistent with


describes in PMID: 19189974 and PMID: 32493277.

Negative control: HepG2 and RD (PMID: 19189974 and PMID:


Anti-Gli1 antibody [EPR4523] (ab134906) at 1/1000 dilution

32493277).

Secondary



Western blot - Anti-Gli1 antibody [EPR4523] (ab134906)




Blocking buffer: 5% NFDM/TBST

Dilution buffer: 5% NFDM/TBST

Predicted band size: 118 kDa Observed band size: 150 kDa

(purified) + RMS13 cell lysate at 10 μg

HRP goat anti-rabbit (H+L) at 1/1000 dilution



Western blot - Anti-Gli1 antibody [EPR4523] (ab134906)

All lanes: Anti-Gli1 antibody [EPR4523] (ab134906) at 1/1000 dilution (unpurified)


Lane 1: RMS13 cell lysate Lane 2: A673 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP-conjugated goat anti-rabbit IgG polyclonal at 1/2000 dilution

Predicted band size: 118 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors