

Anti-Glucocorticoid Receptor antibody ab3671

★★★★☆ [4 Abreviews](#) [2 References](#) [画像数 5](#)

製品の概要

製品名	Anti-Glucocorticoid Receptor antibody
製品の詳細	Rabbit polyclonal to Glucocorticoid Receptor
由来種	Rabbit
特異性	This antibody detects both the unactivated and activated forms of GR.
アプリケーション	適用あり: IHC-P, ICC/IF, WB
種交差性	交差種: Mouse, Human, Snake
免疫原	Synthetic peptide corresponding to Human Glucocorticoid Receptor aa 150-175. Sequence: APTEKEFPKTHSDVSSEQQLKGGTG

 [Run BLAST with](#)

 [Run BLAST with](#)

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.05% Sodium azide Constituent: 99% PBS
精製度	Whole antiserum
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab3671の使用に適用されます**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/250.
ICC/IF	★★★★☆ (1)	1/250.
WB	★★★★☆ (2)	1/500 - 1/2500. Detects a band of approximately 97 kDa (predicted molecular weight: 86 kDa).

ターゲット情報

機能	Receptor for glucocorticoids (GC). Has a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE) and as a modulator of other transcription factors. Affects inflammatory responses, cellular proliferation and differentiation in target tissues. Could act as a coactivator for STAT5-dependent transcription upon growth hormone (GH) stimulation and could reveal an essential role of hepatic GR in the control of body growth. Involved in chromatin remodeling. Plays a significant role in transactivation. Involved in nuclear translocation.
組織特異性	Widely expressed. In the heart, detected in left and right atria, left and right ventricles, aorta, apex, intraventricular septum, and atrioventricular node as well as whole adult and fetal heart.
関連疾患	Defects in NR3C1 are a cause of glucocorticoid resistance (GCRES) [MIM:138040]; also known as cortisol resistance. It is a hypertensive, hyperandrogenic disorder characterized by increased serum cortisol concentrations. Inheritance is autosomal dominant.
配列類似性	Belongs to the nuclear hormone receptor family. NR3 subfamily. Contains 1 nuclear receptor DNA-binding domain.
ドメイン	Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain.
翻訳後修飾	Increased proteasome-mediated degradation in response to glucocorticoids. Phosphorylated in the absence of hormone; becomes hyperphosphorylated in the presence of glucocorticoid. The Ser-203-phosphorylated form is mainly cytoplasmic, and the Ser-211-phosphorylated form is nuclear. Transcriptional activity correlates with the amount of phosphorylation at Ser-211. Sumoylated; this reduces transcription transactivation. Ubiquitinated; restricts glucocorticoid-mediated transcriptional signaling.
細胞内局在	Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand, nuclear after ligand-binding and Nucleus. Localized largely in the nucleus.

画像



Western blot - Anti-Glucocorticoid Receptor antibody (ab3671)

All lanes : Anti-Glucocorticoid Receptor antibody (ab3671) at 1/1000 dilution

Lane 1 : A549 (human lung carcinoma cell line) membrane enriched extract

Lane 2 : MCF7 (human breast adenocarcinoma cell line) membrane enriched extract

Lane 3 : T-47D membrane enriched extract


Lane 4 : MDA-MB-231 (human breast adenocarcinoma cell line) membrane enriched extract

Lane 5 : HeLa (human epithelial cell line from cervix adenocarcinoma) membrane enriched extract

Lane 6 : Mouse brain tissue extract


Lysates/proteins at 30 µg per lane.

Predicted band size: 86 kDa




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor antibody - ChIP Grade (ab3671)

Immunocytochemistry/Immunofluorescence analysis of HEK-293 (Human epithelial cell line from embryonic kidney) cells labeling Glucocorticoid Receptor (green) with ab3671 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor antibody - ChIP Grade (ab3671)

Immunocytochemistry/Immunofluorescence analysis of A2058 (Human metastatic melanoma cell line) cells labeling Glucocorticoid Receptor (green) with ab3671 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.



Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor antibody - ChIP Grade (ab3671)

Immunocytochemistry/Immunofluorescence analysis of HeLa (Human epithelial adenocarcinoma cell line) cells labeling Glucocorticoid Receptor (green) with ab3671 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of *Thamnophis sirtalis* (Common garter snake) brain tissue sections labeling Glucocorticoid Receptor with ab3671 at 1/250. Samples were blocked with 10% goat serum in 0.1M PBS. Samples were incubated with the primary antibody for 48 hours at 4°C. A biotin-conjugated goat anti-rabbit was used as the secondary antibody.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucocorticoid Receptor antibody - ChIP Grade (ab3671)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors