

Anti-Heme Oxygenase 1 antibody [EPR18161-128] ab189491

KO 評価済 リコンビナント RabMAb

★★★★★ [2 Abreviews](#) [39 References](#) [画像数 12](#)

製品の概要

製品名	Anti-Heme Oxygenase 1 antibody [EPR18161-128]
製品の詳細	Rabbit monoclonal [EPR18161-128] to Heme Oxygenase 1
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, IP, ICC/IF, Flow Cyt (Intra)
種交差性	交差種: Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human, mouse and rat spleen lysate. NIH/3T3 and HeLa cell lysate. IHC-P: Human, mouse and rat liver tissue. ICC/IF: HeLa and NIH/3T3 cells. Flow Cyt (intra): HeLa and NIH/3T3 cells. IP: NIH/3T3 whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR18161-128

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab189491の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	1/2000. Predicted molecular weight: 33 kDa.
IHC-P		1/20000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IP		1/30.
ICC/IF	★★★★☆ (1)	1/250.
Flow Cyt (Intra)		1/50.

ターゲット情報

機能

Heme oxygenase cleaves the heme ring at the alpha methene bridge to form biliverdin. Biliverdin is subsequently converted to bilirubin by biliverdin reductase. Under physiological conditions, the activity of heme oxygenase is highest in the spleen, where senescent erythrocytes are sequestered and destroyed.


配列類似性

Belongs to the heme oxygenase family.

細胞内局在

Microsome. Endoplasmic reticulum.

画像



Western blot - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

All lanes : Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491) at 1/2000 dilution

Lane 1 : Wild-type A549 cell lysate

Lane 2 : HMOX1 knockout A549 cell lysate

Lane 3 : Mouse Spleen cell lysate

Lane 4 : HeLa cell lysate

Lane 5 : HEK-293 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution


Performed under reducing conditions.

Predicted band size: 33 kDa

Observed band size: 32 kDa

False colour image of Western blot: Anti-Heme Oxygenase 1 antibody [EPR18161-128] staining at 1/2000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab189491 was shown to bind specifically to Heme Oxygenase 1. A band was observed at 32 kDa in wild-type A549 cell lysates with no signal observed at this size in HMOX1 knockout cell line [ab269503](#) (HMOX knockout A549 lysate [ab259782](#)).


To generate this image, wild-type and HMOX1 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5 % milk in TBS-0.1 % Tween@20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution



Immunohistochemical analysis of paraffin embedded mouse liver tissue labeling Heme Oxygenase 1 with ab189491 at 1/20000 dilution, followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining on Kupffer cells of mouse liver (PMID: 9449694) is observed. Counter stained with hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)



Western blot - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491) at 1/2000 dilution + Human spleen lysate at 10 µg

Secondary

VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) at 1/4000 dilution

Predicted band size: 33 kDa


Observed band size: 28, 32 kDa

Exposure time: 3 minutes

Blocking: 5% NFDM/TBST.

The molecular weight observed is consistent with what has been described in the literature (PMID: 18400743).


The lower band observed is a truncated form of Heme Oxygenase 1 (PMID: 17430897).



Immunocytochemistry/ Immunofluorescence - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Immunofluorescent analysis of 4% paraformaldehyde fixed, 0.1% Triton X-100 permeabilized HeLa (human cervix adenocarcinoma epithelial cell) cells labeling Heme Oxygenase 1 with ab189491 at 1/250 dilution, followed by [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on HeLa cells. Details of counterstains: [ab195889](#) Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at a 1/200 dilution; DAPI for nuclei.


The negative controls are as follows: Secondary antibody only for control.



Immunocytochemistry/ Immunofluorescence - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Immunofluorescent analysis of 4% paraformaldehyde fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labeling Heme Oxygenase 1 with ab189491 at 1/250 dilution, followed by **ab150077** AlexaFluor[®]488 Goat anti-Rabbit secondary at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on NIH/3T3 cells. Details of counterstains: **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594) at a 1/200 dilution; DAPI for nuclei.

The negative controls are as follows: Secondary antibody only for control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Immunohistochemical analysis of paraffin embedded human liver tissue labeling Heme Oxygenase 1 with ab189491 at 1/20,000 dilution, followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining on Kupffer cells of human liver (PMID: 9449694) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Perform heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Immunohistochemical analysis of paraffin embedded rat liver tissue labeling Heme Oxygenase 1 with ab189491 at 1/20,000 dilution, followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining on Kupffer cells of rat liver (PMID: 9449694) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).



Western blot - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

All lanes : Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491) at 1/5000 dilution

Lane 1 : NIH/3T3 (mouse embryonic fibroblast) lysate

Lane 2 : Rat spleen lysate

Lane 3 : HeLa (human cervix adenocarcinoma epithelial cell) lysate

Lane 4 : Mouse spleen lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 33 kDa

Observed band size: 28, 32 kDa

Blocking: 5% NFDm/TBST.


Exposure time:

Lanes 1,2 and 3: 2 seconds;

Lane 4: 1 second


The molecular weight observed is consistent with what has been described in the literature (PMID: 18400743).

The lower band observed is a truncated form of Heme Oxygenase 1 (PMID: 17430897).




Flow Cytometry (Intracellular) - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Intracellular flow cytometric analysis of 90% methanol/ 4% paraformaldehyde fixed NIH/3T3 (mouse embryonic fibroblast) cell line labeling Heme Oxygenase1 with ab189491 at 1/50 (Red) compared with a Rabbit monoclonal IgG (**ab172730**) (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**), at 1/2000 dilution was used as the secondary antibody.



Flow Cytometry (Intracellular) - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Intracellular flow cytometric analysis of 90% methanol/4% paraformaldehyde fixed HeLa (human cervix adenocarcinoma epithelial cell) cell line labeling Heme Oxygenase1 with ab189491 at 1/50 (Red) compared with a Rabbit monoclonal IgG (**ab172730**) (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**), at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Heme Oxygenase 1 was immunoprecipitated from 0.35 mg of NIH/3T3 (mouse embryonic fibroblast) whole cell lysate with ab189491 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab189491 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)), was used for detection at 1/10,000 dilution.

Lane 1: NIH/3T3 (mouse embryonic fibroblast) whole cell lysate 10 µg (Input).

Lane 2: NIH/3T3 whole cell lysate (+).

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab189491 in NIH/3T3 whole cell lysate (-).

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

The truncated form of Heme Oxygenase 1 is described in the literature (PMID: 17430897).

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab189491)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors