

Anti-HHV8 antibody [LN53] ab4103

14 References [画像数 1](#)

製品の概要

製品名	Anti-HHV8 antibody [LN53]
製品の詳細	Rat monoclonal [LN53] to HHV8
由来種	Rat
特異性	HHV 8 is the likely etiological agent of Kaposi's sarcoma (KS). HHV 8 encodes a latent nuclear antigen (LNA), which is the product of the viral gene orf 73. LNA is capable of forming a complex with retinoblastoma susceptibility gene product, which may be related to its oncogenic activity.
アプリケーション	適用あり: IHC-P, IHC-Fr
種交差性	交差種: Human herpesvirus 8
免疫原	Recombinant full length protein corresponding to HHV8. Corresponding to the latent nuclear antigen 1 molecule of HHV 8.
エピトープ	Located within the ORF-73 region.
ポジティブ・コントロール	Kaposi's sarcoma.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.3</p> <p>Preservative: 0.05% Sodium azide</p> <p>Constituents: Tissue culture supernatant, 1% BSA</p>
精製度	IgG fraction
ポリ/モノ	モノクローナル

クローン名	LN53
アイソタイプ	IgG2c

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab4103の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/25 - 1/50. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. ABC method.
IHC-Fr		1/25 - 1/50. ABC method.

ターゲット情報

関連性 HHV8 has been found to be associated with three different diseases observed in AIDS patients; kaposi's sarcoma, primary effusion lymphoma (which is a rare type of non-Hodgkin lymphoma affecting the body cavities) and multicentric Castleman's disease. To date there is much evidence to support a direct role for HHV8 in kaposi's sarcoma.

画像



Cultured cells infected with HHV8, stained with HHV8 using ABC and DAB chromogen.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HHV8 antibody [LN53] (ab4103)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors