

Anti-HMGB2 antibody [EPR6302] ab133540

KO 評価済 リコンビナント RabMAb


2 References 画像数 7

製品の概要

製品名	Anti-HMGB2 antibody [EPR6302]
製品の詳細	Rabbit monoclonal [EPR6302] to HMGB2
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P 適用なし: ICC/IF or IP
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HEK-293T, HAP1, K562, HL-60, C6, RAW 264.7, and PC-12 cell lysates. IHC-P: Human breast tissue and Human colon carcinoma tissue.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
解離定数 (K _D 値)	K _D = 2.97 x 10 ⁻¹⁰ M



[Learn more about K_D](#)

バッファー	pH: 7.2 Preservative: 0.05% Sodium azide
-------	---

	Constituents: 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR6302
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab133540の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/10000. Predicted molecular weight: 24 kDa.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

追加情報 Is unsuitable for ICC/IF or IP.

ターゲット情報

機能	DNA binding proteins that associates with chromatin and has the ability to bend DNA. Binds preferentially single-stranded DNA. Involved in V(D)J recombination by acting as a cofactor of the RAG complex. Acts by stimulating cleavage and RAG protein binding at the 23 bp spacer of conserved recombination signal sequences (RSS).
配列類似性	Belongs to the HMGB family. Contains 2 HMG box DNA-binding domains.
細胞内局在	Nucleus. Chromosome.

画像



Western blot - Anti-HMGB2 antibody [EPR6302]
(ab133540)

All lanes : Anti-HMGB2 antibody [EPR6302] (ab133540) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : HMGB2 knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 24 kDa

Observed band size: 24 kDa

Lanes 1- 2: Merged signal (red and green). Green - ab133540 observed at 24 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) observed at 37 kDa.


ab133540 was shown to react with HMGB2 in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line **ab266358** (knockout cell lysate **ab257156**) was used. Wild-type HEK-293T and HMGB2 knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab133540 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HMGB2 antibody
[EPR6302] (ab133540)

Immunohistochemistry analysis of paraffin embedded Human breast tissue labelling HMGB2 with ab133540 at 1/250 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-HMGB2 antibody [EPR6302] (ab133540)

All lanes : Anti-HMGB2 antibody [EPR6302] (ab133540) at 1/1000 dilution

Lane 1 : Wild-type HAP1 whole cell lysate

Lane 2 : HMGB2 knockout HAP1 whole cell lysate

Lane 3 : HeLa cell lysate

Lane 4 : K562 cell lysate


Lysates/proteins at 20 µg per lane.

Predicted band size: 24 kDa

Observed band size: 24 kDa

Lanes 1 -4: Merged signal (red and green). Green - ab133540 observed at 24 kDa. Red - loading control, **ab9484**, observed at 37 kDa.


ab133540 was shown to recognize HMGB2 in wild-type HAP1 cells as signal was lost at the expected MW in HMGB2 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and HMGB2 knockout samples were subjected to SDS-PAGE. Ab133540 and **ab9484** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HMGB2 antibody [EPR6302] (ab133540)

Immunohistochemistry analysis of paraffin embedded Human colon carcinoma tissue labelling HMGB2 with ab133540 at 1/250 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-HMGB2 antibody [EPR6302]
(ab133540)

All lanes : Anti-HMGB2 antibody [EPR6302] (ab133540) at 1/1000 dilution

Lane 1 : K562 cell lysate

Lane 2 : HL-60 cell lysate


Lane 3 : C6 cell lysate

Lane 4 : RAW 264.7 cell lysate

Lane 5 : PC-12 cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 24 kDa



SPR Scanning - Anti-HMGB2 antibody [EPR6302]
(ab133540)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-HMGB2 antibody [EPR6302] (ab133540)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors