# abcam

## Product datasheet

## Anti-hnRNP A1 (citrulline R140) antibody [EPR20178] - BSA and Azide free ab251488

יעלאעבע RabMAb

#### 画像数3

#### 製品の概要

特記事項

製品名 Anti-hnRNP A1 (citrulline R140) antibody [EPR20178] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EPR20178] to hnRNP A1 (citrulline R140) - BSA and Azide free

由来種 Rabbit

特異性 Based on sequence homology, we predict that this antibody may react with HNRNPA1L2.

アプリケーション 適用あり: Dot blot, WB

種交差性 交差種: Recombinant fragment

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ab251488 is the carrier-free version of ab208030.

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb patents**.

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

**バッファー** pH: 7.2

Constituent: PBS

キャリア・フリー はい

**ポリ/モノ** モノクローナル **ウローン名** EPR20178

アイソタイプ IgG

#### アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab251488の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項  |
|----------|-----------|---|
| Dot blot |           | Use at an assay dependent concentration.  |
| WB       |           | Use at an assay dependent concentration. Detects a band of approximately 34 kDa (predicted molecular weight: 39 kDa). |

#### ターゲット情報

機能 Involved in the packaging of pre-mRNA into hnRNP particles, transport of poly(A) mRNA from the

nucleus to the cytoplasm and may modulate splice site selection. May play a role in HCV RNA

replication.

**配列類似性** Contains 2 RRM (RNA recognition motif) domains.

翻訳後修飾 Arg-194, Arg-206 and Arg-225 are dimethylated, probably to asymmetric dimethylarginine.


Sumoylated.

**細胞内局在** Nucleus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs.

Shuttles continuously between the nucleus and the cytoplasm along with mRNA. Component of ribonucleosomes. In the course of viral infection, colocalizes with HCV NS5B at speckles in the

cytoplasm in a HCV-replication dependent manner.


### 画像



Dot Blot - Anti-hnRNP A1 (citrulline R140) antibody [EPR20178] - BSA and Azide free (ab251488)

This data was developed using <u>ab208030</u>, the same antibody clone in a different buffer formulation.Dot blot analysis of hnRNP A1 (citrulline R140) labeled with <u>ab208030</u> at 1/1000 dilution. **Lane 1:** hnRNP A1 (citrulline R140) peptide. **Lane 2:** hnRNP A1 (citrulline R140) noncitrulline peptide.

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution was used as secondary antibody. Blocking and dilution buffer: 5% NFDM/TBST. Exposure time: 3 minutes.



Western blot - Anti-hnRNP A1 (citrulline R140) antibody [EPR20178] - BSA and Azide free (ab251488)

**All lanes**: Anti-hnRNP A1 (citrulline R140) antibody [EPR20178] (ab208030) at 1/1000 dilution

**Lane 1 :** Empty vector (vector control) transfected HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

**Lane 2**: HEK-293T whole cell lysate transfected with GFP tagged PADI2 and treated with 10mM CaCI2 for 2h, then 10uM lonomycin for 2h

**Lane 3**: Empty vector (vector control) transfected HEK-293T whole cell lysate

**Lane 4**: HEK-293T whole cell lysate transfected with GFP tagged PADI4 and treated with 10mM CaCI2 for 2h, then 10uM lonomycin for 2h

Lysates/proteins at 10 µg per lane.

#### **Secondary**

**All lanes :** Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/100000 dilution


Developed using the ECL technique.

**Predicted band size:** 39 kDa **Observed band size:** 34 kDa

Exposure time: 5 seconds

This data was developed using <u>ab208030</u>, the same antibody clone in a different buffer formulation.

Blocking and dilution buffer: 5% NFDM/TBST.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors