


Anti-hnRNP H antibody ab10374

★★★★★ **5 Abreviews** **29 References** 画像数 3

製品の概要

製品名	Anti-hnRNP H antibody
製品の詳細	Rabbit polyclonal to hnRNP H
由来種	Rabbit
特異性	The antibody was designed against hnRNPH1 and does not recognize hnRNPH3. The epitope recognized by ab10374 maps to a region between residue 400 and the C-terminus (residue 448) of human Heterogeneous Nuclear Ribonucleoprotein H1 using the numbering given in entry NP_005511.1 (GeneID 3187). This product may cross-react with hnRNPH2.
アプリケーション	適用あり: WB, IHC-P, IP, ICC/IF, IHC-Fr
種交差性	交差種: Mouse, Human 交差が予測される動物種: Rat, Rabbit, Horse, Guinea pig, Dog, Pig, Chimpanzee, Rhesus monkey, Gorilla, Opossum, Orangutan, Elephant 
免疫原	Synthetic peptide corresponding to Human hnRNP H aa 400 to the C-terminus (C terminal). Database link: 3187
ポジティブ・コントロール	IP: HeLa whole cell lysate WB: HeLa, HEK-293T, and NIH/3T3 cell lysates.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7 Preservative: 0.1% Sodium azide Constituents: 0.021% PBS, 1.764% Sodium citrate, 1.815% Tris
精製度	Immunogen affinity purified
特記事項 (精製)	Antibodies were affinity purified using the peptide immobilized on solid support.

ポリ/モノ ポリクローナル
アイソタイプ IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab10374の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (2)	1/10000 - 1/30000. Detects a band of approximately 49 kDa (predicted molecular weight: 53 kDa).
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
IP	★★★★★ (1)	Use at 4-10 µg/mg of lysate.
ICC/IF	★★★★★ (2)	Use at an assay dependent concentration. PubMed: 20020773
IHC-Fr		Use at an assay dependent concentration. PubMed: 24462217

ターゲット情報

機能	This protein is a component of the heterogeneous nuclear ribonucleoprotein (hnRNP) complexes which provide the substrate for the processing events that pre-mRNAs undergo before becoming functional, translatable mRNAs in the cytoplasm. Mediates pre-mRNA alternative splicing regulation. Inhibits, together with CUGBP1, insulin receptor (IR) pre-mRNA exon 11 inclusion in myoblast. Binds to the IR RNA. Binds poly(RG).
組織特異性	Expressed ubiquitously.
配列類似性	Contains 3 RRM (RNA recognition motif) domains.
ドメイン	Each quasi-RRM repeat bound poly(RG), while only the N-terminal QRRM bound poly(RC) and poly(RU). None of the repeats bound detectable amounts of poly(RA).
細胞内局在	Nucleus > nucleoplasm.

画像



All lanes : Anti-hnRNP H antibody (ab10374) at 0.2 µg/ml

Lane 1 : HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 50 µg

Lane 2 : HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 15 µg

Lane 3 : HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 5 µg


Lane 4 : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate at 50 µg

Lane 5 : NIH/3T3 (mouse embryo fibroblast cell line) whole cell lysate at 50 µg


Developed using the ECL technique.

Predicted band size: 53 kDa

Exposure time: 30 seconds



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human ovarian carcinoma tissue labelling hnRNP H with ab10374 at 1/1000 (1µg/ml). Detection: DAB.



Detection of human hnRNP H by immunoprecipitation of whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells.

Lane 1: Rabbit polyclonal to hnRNP H (ab10374) at 0.2 µg/mg lysate current batch

Lane 2: Rabbit polyclonal to hnRNP H (ab10374) at 0.2 µg/mg lysate former batch.

Lane 3: Control IgG

ab10374 was used at 1 µg/mL for Western blot.

Detection: Chemiluminescence with an exposure time of 30 seconds.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors