

HRP Anti-6X His tag® antibody [AD1.1.10] ab178563

2 References [画像数 1](#)

医薬用外劇物

製品の概要

製品名	HRP Anti-6X His tag® antibody [AD1.1.10]
製品の詳細	HRP Mouse monoclonal [AD1.1.10] to 6X His tag®
由来種	Mouse
標識	HRP
アプリケーション	適用あり: WB, ELISA
種交差性	交差種: Species independent
免疫原	Synthetic peptide corresponding to 6X His tag®. Antibody was raised against a polypeptide containing a tag
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Lyophilized:Reconstitute to 1.0 ml with distilled deionized water vortex gently and allow vial to sit on ice for 20 minutes; concentration will be 0.1mg/ml. We recommend that the investigator determine the appropriate working concentration for their specific application. Avoid exposure to heat and light.
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze. Store In the Dark.
バッファー	pH: 7.40 Preservative: 0.01% Thimerosal (merthiolate) Constituents: 1.4% Sodium phosphate, 0.58% Sodium chloride, 1.5% BSA, 1.71% Sucrose Buffer after reconstitution in 1.0 ml distilled deionized water.
精製度	Size exclusion
ポリ/モノ	モノクローナル

クローン名 AD1.1.10
アイソタイプ IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab178563の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.

ターゲット情報

関連性 The H-H-H-H-H motif is used as a tag on many recombinant proteins to facilitate purification. His-tags can be fused to the amino- or carboxy- termini of proteins in transfected or transformed cells.

細胞内局在 Depends upon the localization of the parent protein tagged with hexahistidine.

画像



Western Blot of a 41kDa protein fused (expressed in E.Coli) to His tag using ab178563 0.25 µg/ml. Loading was confirmed by Coomassie blue staining of transferred proteins.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors