abcam

Product datasheet

HRP Anti-HDAC3 antibody [Y415] ab208356

יעלאעבע RabMAb

画像数 2

製品の概要

製品名 HRP Anti-HDAC3 antibody [Y415]

製品の詳細 HRP Rabbit monoclonal [Y415] to HDAC3

由来種 Rabbit 標識 HRP

アプリケーション **適用あり:** WB 種交差性 交差種: Human

交差が予測される動物種: Mouse. Rat 4

免疫原 Synthetic peptide within Human HDAC3 aa 400 to the C-terminus. The exact immunogen

> sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact

our Scientific Support team to discuss your requirements.

Database link: O15379

Run BLAST with Run BLAST with

ポジティブ・コントロール WB: K562 whole cell lysate.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 特記事項

monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

Protein A purified 精製度

ポリモノ モノクローナル

クローン名 Y415

アイソタイプ lgG

アプリケーション

Abpromise保証は、次のテスト済みアプリケーションにおけるab208356の使用に適用されます The Abpromise guarantee アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/2000. Detects a band of approximately 49 kDa (predicted molecular weight: 49 kDa).

ターゲット情報

機能

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation.

組織特異性 Widely expressed.


配列類似性 Belongs to the histone deacetylase family. HD type 1 subfamily.

翻訳後修飾 Sumoylated in vitro.

細胞内局在 Nucleus.

画像

(ab208356)



HRP Anti-HDAC3 antibody [Y415] (ab208356) at 1/2000 dilution + K562 (Human erythromyeloblastoid leukemia cell line) Whole Cell Lysate at 10 µg

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 49 kDa Observed band size: 49 kDa

Exposure time: 20 minutes

buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% milk before being incubated with ab208356 overnight at 4°C. Antibody binding was visualised using ECL development solution **ab133406**.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors